

Sensorimotor Training for Shoulder Injury

Literature Review

Joseph B. Myers, PhD, ATC; and Sakiko Oyama, MS, ATC

■ ABSTRACT

When the shoulder is injured, a cascade of interrelated events result, including tissue pathology, pain, and sensorimotor alterations. Clinicians use sensorimotor training to address these events. The purpose of this review was to discuss how the sensorimotor system contributes to shoulder function and to review evidence regarding the effectiveness of training to target sensorimotor mechanisms. Review of peer-reviewed literature was performed with both Medline and Cumulative Index in Nursing and Health Literature databases. For studies that involved some component of shoulder sensorimotor training, only those published within the past 10 years were included. The search yielded 23 publications of varying levels of evidence, suggesting the ability of sensorimotor training to result in both acute and chronic adaptations to the sensorimotor system. Although the evidence regarding usage still needs to improve, some data suggest that the sensorimotor system may be trainable.

It is commonly accepted that the joints of the shoulder complex rely heavily on coordinated activation of the muscles that cross each joint for much needed stability and performance of functional activity. The sensorimotor system is primarily responsible for regulating this coordinated muscle activation of the shoulder

musculature.¹ When injury to the shoulder occurs, not only do tissue pathology and pain result, but potential alterations to the sensorimotor system can also manifest. As such, clinicians who rehabilitate the shoulder not only have to address the tissue pathology and pain, but also must retrain the sensorimotor mechanisms that are important to shoulder stability and function.

The purpose of this review is to discuss how the sensorimotor system contributes to shoulder function, describe how it is affected with injury, and review evidence regarding the effectiveness of sensorimotor training to target the mechanisms necessary for stability and function of shoulder complex. To complete this review, Internet-based searches of the peer-reviewed literature were performed with both Medline and Cumulative Index in Nursing and Health Literature (CINAHL) databases. Searches were performed with the following keywords (and combinations of keywords): *shoulder, scapula, sensorimotor, proprioception, neuromuscular, muscle activation, electromyography, rehabilitation, and training*. For the review of evidence, the titles and abstracts of 683 articles were reviewed to identify orthopedic-related articles that either measured change in the sensorimotor system that resulted from some training intervention or measured some sensorimotor characteristic during performance of a sensorimotor training activity. Only studies published within the past 10 years were reviewed. For the intervention studies reviewed, only those that met criteria for inclusion in the Physiotherapy Evidence Database (PEDro) were included. For an intervention to be included in this review (ie, meet PEDro inclusion criteria), the study had to:

- Compare at least 2 interventions (intervention versus control or sham, or 2 interventions).
- Include interventions that are currently part of clinical

The authors are from the Department of Exercise and Sport Science, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina.

Originally submitted February 4, 2009.

Accepted for publication May 14, 2009.

The authors have no financial or proprietary interest in the materials presented herein.

Address correspondence to Joseph B. Myers, PhD, ATC, Department of Exercise and Sport Science, University of North Carolina at Chapel Hill, CB#8700 Fetzer, Chapel Hill, NC 27599-8700; e-mail: joemyers@email.unc.edu.

doi:10.3928/19425864-2009xxxx-xx

practice or could become part of clinical practice.

- Include interventions applied to human subjects who are representative of those to whom the intervention might be applied in the course of clinical practice.
- Involve random allocation or intended-to-be-random allocation of participants to interventions.
- Be published as a full paper in a peer-reviewed journal.

SENSORIMOTOR CONTRIBUTION AT THE SHOULDER

The sensorimotor system includes sensory, motor, and central integration and processing components involved with maintenance of joint homeostasis during functional activity.² The role of the sensorimotor system is to modulate ongoing muscle activation characteristics by using afferent neural input to initiate and adjust efferent neural signal to the desired muscles. This efferent response to a constantly changing afferent signal occurring unconsciously for the purpose of maintaining joint stability is called *neuromuscular control*.² Proprioception is one of the afferent neural inputs from the periphery used by the sensorimotor system, conveying information regarding joint movement (kinesthesia), position and orientation, and force applied to or present within a joint.³ During joint movement, tension and force develop within the tissues (muscles, tendons, fascia, ligaments, and joint capsule) about the shoulder. The result is stimulation of mechanoreceptors, which translates the mechanical change in the tissue (length and tension changes) as a neural signal to the central nervous system via afferent neural pathways. Within the central nervous system, the proprioception information regarding length and tension is integrated with the other peripheral senses, such as tactile sensation, vision, vestibular input, and descending neural signals from spinal cord, brain stem, cerebellum, and cortex.⁴ The result is neural signals conveyed through efferent pathways to the motor neurons to produce muscle activation or inhibition and to regulate sensitivity of the mechanoreceptors within the muscle (muscle spindle).⁵ Regulation of the sensitivity of muscle spindle through the efferent neural signal, in turn, influences the afferent neural signals sent to the central nervous system.⁵

Both static joint stability (stability provided by osseous geometry, capsuloligamentous and tenomuscular restraints, and intra-articular pressure) and dynamic joint stability (stability provided by the contractile action by muscles) contribute to functional joint stability. How-

ever, the contribution of the dynamic stabilizers outweighs the contribution from the static stabilizers at the glenohumeral joint because of poor glenohumeral joint congruency and laxity of the glenohumeral joint capsule that permit large range of motion.⁶⁻⁸ The sensorimotor system serves a critical role in providing dynamic joint stability through its control over the activation of muscles that cross the shoulder joint.^{4,9} Modulation of tone (and ultimately stiffness) of the muscles surrounding the shoulder by regulation of muscle spindle sensitivity is most likely how the sensorimotor system provides joint stability at the shoulder.¹⁰ When the muscle spindle is highly sensitive, relatively small changes in muscle length and tension triggers afferent neural signals, which results in enhanced muscle tone (muscle activation level) and shorter latency period of the muscle reflexive responses.¹¹ Increased muscle tone and shorter reflex latency period provide stiffness to the muscle-tendon unit, which makes the shoulder more resistant to perturbing events of the humeral head or scapula. In a recent study, moderate level of muscle activation was shown to result in significant increase in glenohumeral joint stiffness,¹² demonstrating the vital role that muscle activation plays in increasing the joint stiffness and aiding joint stability at the glenohumeral joint. Increased muscle stiffness also improves joint stability by providing tension on the joint capsule, a mechanism referred to as *capsular dynamitization*.¹³ The tendons of the rotator cuff muscles and the other muscles that attach onto the proximal humerus (pectoralis major, latissimus dorsi, and teres major) blend into the joint capsule.¹⁴ As result, the glenohumeral joint capsule tightens and stiffens when these muscles are activated.

In addition to regulation of the joint stiffness, sensorimotor system contributes to joint stability by coordinating activities of the muscles surrounding the joint. Co-activation of the rotator cuff muscles is particularly important in producing joint compression to keep the humeral head centered within the glenoid fossa.¹⁵⁻²¹ When the 4 rotator cuff muscles co-activate, the muscles collectively produce a resultant joint compressive force, centralizing the humeral head within the glenoid, minimizing translation of the humeral head.¹⁵⁻²² For example, anterior translation of the humeral head increased by approximately half [**Query #1: Can “half” be substituted by “50%” for consistency with the latter part of this sentence?**], when the compression force by the rotator cuff muscles decreased by 50%.²³ In addition to the compressive force, co-activation of the deltoid and the rotator

cuff muscles allows rotation of the humeral head on glenoid fossa while minimizing superior translation of the humerus during humeral elevation.^{24,25}

In addition to the muscles about the glenohumeral joint, precise activation of the muscles controlling scapular movement and stability are also necessary for shoulder stability and function. For example, Moseley et al²⁶ examined the electromyographic activity of the 3 portions of the trapezius, levator scapula, rhomboids, pectoralis minor, and serratus anterior during different scapular training exercises and noted that majority of the muscles were active during all movement, suggesting that the scapular muscles play important roles during scapular movement. During humeral elevation, the electromyography literature reporting scapular muscle activation suggest that the upper trapezius, lower trapezius, and serratus anterior work in synergy to achieve the upward scapular rotation necessary for humeral elevation to occur.²⁶⁻³¹ Efficient scapular movement is important in positioning and orienting the glenoid fossa in alignment with the humerus so that the line of pull and the length tension relationship of the rotator cuff muscles can be maintained.³²

Effect of Injury on Sensorimotor System

Given the important roles the sensorimotor system plays on shoulder stability and function, sensorimotor deficits associated with injury are of great concern clinically and should be addressed following injury. From a review of the literature, it appears that the sensorimotor system can be affected by joint or muscle injuries that cause pain, trauma, or attenuation of the capsuloligamentous and musculotendinous tissues.

The use of experimental pain has allowed the investigators to isolate the effect of pain in absence of mechanical damage to the tissue or the mechanoreceptors.³³⁻³⁸ Research using experimental pain models have demonstrated changes in sensorimotor system activity simply due to pain.³³⁻³⁸ Decreased excitability of the cortical and spinal motor neurons in presence of experimental pain demonstrated by one of the studies³⁸ suggests the existence of central mechanism (ie, inhibition initiated by the motor cortex), resulting in the sensorimotor system alteration. Research has also demonstrated alteration in muscle activation,^{33,34,37,39} motor unit firing rate,³⁵ and maximal force production³⁶ when pain is present. Alternatively, improvements in muscle activation and scapular movement pattern have also been reported in patients with rotator cuff disease who received anesthetic injec-

tion.^{40,41}

In addition to the pain that occurs with injury, attenuation and stretching of the capsuloligamentous and musculotendinous structures may also contribute to the sensorimotor function deficits. Alterations in the sensorimotor system, including deficit in proprioception,⁴²⁻⁴⁴ altered muscle activity during upper extremity tasks,^{23,45-48} altered reflexive muscle activity,⁴⁹ and dysfunctional movement patterns⁵⁰ have been reported in patients with anterior and multidirectional instability. Alteration in the proprioception conveyed to the central nervous system due to reorganization of the length and tension appreciation by the mechanoreceptors and deafferentation of the mechanoreceptors within the tissue or the neural pathways are considered the primary mechanisms for these sensorimotor deficits.^{51,52} Because patients with joint instability are typically affected by both joint hyperlaxity and pain,^{45,53-55} it is impossible to isolate the effect of joint laxity from that of pain. However, other findings demonstrating decreased sensorimotor function in individuals with asymptomatic joint laxity suggest the laxity itself attributes to sensorimotor deficits in individuals with instability.⁵⁶⁻⁵⁸ Studies reporting restoration of sensorimotor function in patients with shoulder instability following a surgery to address capsule laxity may also support the influence of joint laxity on sensorimotor system.^{42,44,59,60} However, the accompanying decrease in pain that most likely resulted from surgery or the treatment following surgery cannot be overlooked.

Sensorimotor dysfunction has also been demonstrated in patients with varying degrees of rotator cuff tendinopathy (subacromial impingement to rotator cuff tears). The dysfunction found in these patients includes decreased ability to maintain steady force production⁶¹; decreased activity of the middle deltoid during moderately strenuous external rotation task⁶²; increased activation of the biceps brachii^{63,64}; alteration of the muscle activity ratio among the upper, middle, and lower portions of the trapezius muscles⁴⁶; disrupted co-activation between the middle trapezius and serratus anterior muscles³⁵; suppressed activation and coactivation of the rotator cuff muscles^{25,65}; and delayed onset of the muscles around the scapula during elevation tasks.⁶⁶ Currently, it is unknown whether these sensorimotor alterations contributed to or result from the tendinopathy.

Although the specific patterns of sensorimotor alterations observed in the studies mentioned above may vary, it is evident that the sensorimotor system is affected by

TABLE 1

Review Summary of Acute Adaptation (EMG Research Findings) from Shoulder Sensorimotor Training (1998 to Present)^a

STUDY	CONTROL PARTICIPANTS		PARTICIPANTS WITH INJURY		TRAINING USED	PERTINENT FINDINGS
Decker et al ²⁷	N		N		OKC, CKC	Identified effective exercises to stimulate the serratus anterior
Decker et al ⁹⁴	N		N		OKC, CKC	Identified exercises that best stimulate each portion of the subscapularis
Ekstrom et al ²⁹	N		N		OKC	Identified which exercises maximally activated the trapezius and serratus anterior muscles
Hintermeister et al ⁹⁵	N		N		OKC	Identified exercise that best targets the rotator cuff
Kibler et al ⁹³	Y		Y		OKC	No difference in muscle activation between symptomatic and asymptomatic individual when performing exercise
Kinney et al ¹⁰¹	N		N		OKC	Demonstrated which position of humeral abduction best isolates the middle and lower portion of the trapezius muscle during prone horizontal abduction exercises
Lehman et al ⁹⁸	N		N		CKC	Training on an unstable surface (Swiss ball) had no effect on activation of shoulder girdle muscles
Lehman et al ⁹⁹	N		N		CKC	Training on an unstable surface (Swiss ball) increased muscle activation
Lister et al ¹⁰⁰	N		N		OKC, OD	Training using an oscillatory device resulted in greater scapular stabilizer activation
Ludewig et al ³⁰	Y		Y		CKC	No difference in serratus anterior and trapezius muscle activation between symptomatic and asymptomatic individuals when performing CKC exercise
Myers et al ³¹	N		N		OKC	Demonstrated which throwing related rubber tubing exercises result in the most activation of all muscles of the shoulder girdle
Reinold et al ¹⁰²	N		N		OKC	Demonstrated which "full can" and "empty can" exercises best isolate the supraspinatus and deltoid (middle and posterior) muscles
Reinold et al ¹⁰³	N		N		OKC	Demonstrated which external humeral rotation strengthening exercises best activated the rotator cuff and deltoid muscles
Uhl et al ⁹⁷	N		N		CKC	Exercises that involved progressively increased upper extremity weight bearing resulted in increased shoulder girdle muscle activation

Abbreviations: OKC, open kinetic chain resistance exercises (cuff weights and elastic bands); CKC, closed kinetic chain exercises performed in an upper extremity weight bearing position; OD, exercises that involve oscillatory devices (ex. Bodyblade).

^a None of these articles qualified for a PEDro score assignment

injury and pain. In addition, it seems reasonable to believe that changes in the central nervous system, as well as damage to the peripheral structures, contribute to the alteration in sensorimotor system following injury, although the exact neurophysiologic mechanism is still unclear. However, it is important to acknowledge that the sensorimotor function has never been assessed prospectively, and therefore, existence of sensorimotor deficit prior to

the occurrence of injury cannot be ignored. Particularly for the shoulder injuries that are chronic in nature, such as rotator cuff tears, chronic instability, or subacromial impingement, it is possible that the sensorimotor deficit may have existed prior to the injury and contributed to the occurrence of the injury. This is a weakness in the current knowledge base regarding risk factors to shoulder injuries. Nonetheless, the studies demonstrating the

TABLE 2

Review Summary of Chronic Adaptation from Shoulder Sensorimotor Training (1998 to Present)

STUDY	PEDro SCORE	TRAINING USED	VARIABLES MEASURED	PERTINENT FINDINGS
Carter et al ¹⁰⁴	5/10	Plyo	Strength & Throwing velocity	Plyometrics resulted in increased concentric and eccentric strength of the humeral rotators and throwing velocity
Ginn and Cohen ¹¹¹	6/10	CR	Strength, ROM, Scapular movement, Pain, & Function	Demonstrated that a clinical rehabilitation program that includes sensorimotor training is an effective conservative treatment option for shoulder pain
Kraemer et al ¹¹⁴	6/10	OKC	Strength & Power output, and Function (Tennis serve velocity)	Demonstrated that sport specific resistance training that utilizes multiple sets results in improved strength, power output, and tennis serve velocity in women tennis players
Padua et al ¹⁰⁸	6/10	OKC, CKC, PNF	Strength, Prop, & Functional movement	Shoulder strength improved following 5 weeks of OKC and PNF exercise. Functional throwing performance improved with PNF
Rogol et al ¹⁰⁷	5/10	OKC, CKC	Prop	Both OKC and CKC exercises resulted in improved proprioception
Schulte-Edelmann et al ¹⁰⁵	5/10	Plyo	Strength	A plyometric program resulted in increased power output by the elbow extensor muscles
Swanik et al ¹⁰⁶	6/10	Plyo	Prop & Strength	Plyometrics resulted in both proprioception and improved strength characteristics
Swanik et al ¹¹²	6/10	OKC, CKC	Strength & Pain	A six week function training program resulted in decreased incidence of shoulder pain in collegiate swimmers, but no strength gains resulted
Ubinger et al ¹¹⁰	5/10	CKC	WBS	A CKC training program resulted in improved shoulder stability in a weight bearing position

Abbreviations: OKC, open kinetic chain resistance exercises (cuff weights, elastic bands); CKC, closed kinetic chain exercises performed in an upper extremity weight bearing position; Prop, proprioception; Plyo, plyometric exercises; CR, clinical rehabilitation program that involves stretching and various forms of strengthening; PNF, proprioceptive neuromuscular facilitation exercises; WBS, stability when the upper extremity is placed in a weight bearing position; ROM, range of motion.

sensorimotor deficit following injuries discussed thus far clearly indicate the need for training the sensorimotor system following injury.

Compounding the influence of injury and pain is the effect of fatigue on the sensorimotor system. Fatigue may also contribute to the sensorimotor deficits demonstrated with pain and injury. Components of the sensorimotor system, including proprioception,⁶⁷⁻⁷⁵ muscle activation characteristics,⁷⁶⁻⁸¹ and movement characteristics and coordination,^{76,82-92} appear to be adversely affected by fatigue. The susceptibility of the sensorimotor system to fatigue effects suggests the need for training to improve resistance to fatigability, ultimately decreasing the alterations that manifest with the onset of fatigue.

EFFICACY OF SHOULDER SENSORIMOTOR TRAINING

Given the important role of sensorimotor mechanisms in shoulder stability and function, clinicians use a variety of training techniques in hopes of restoring these mechanisms that are altered with injury. Sensorimotor training activities are believed to improve activation characteristics and force production by the muscles about the shoulder girdle, alter motor programs, and improve proprioceptive acuity. Specifically, they are believed to stimulate activation and coordination of muscles that might be inhibited or altered following injury, facilitate muscle coactivation vital to the force couple mechanisms, and enhance the reflexive loops associated with joint stability. In addition, it is hoped that by improving the activation characteristics, the proprioceptive feedback from the musculature might be improved as well, through enhanced muscle spindle performance

and muscle stiffness. The most common sensorimotor training techniques described in the literature we reviewed included activities that incorporate joint position and movement repositioning, performance of exercise in close kinetic chain positions, proprioceptive neuromuscular facilitation, plyometrics, joint perturbations, selective activation of specific muscles, or global activation of many muscles about the shoulder girdle. The literature includes research studies that demonstrate acute adaptation to the sensorimotor system from performing specific exercises (Table 1) and chronic adaptation to the sensorimotor system from performing training exercises over a period of time (Table 2).

In our review of the literature, we found 14 studies published in the past 10 years that used electromyography to measure the acute adaptation (activation characteristics) that occurs while performing therapeutic exercises (Table 1). Because these studies do not use intervention research designs that meet PEDro requirements, no score was assigned. Yet the results of these studies do provide some valuable information for clinicians, which warrants a discussion. From the measured activation characteristics, recommendations are often made regarding which exercises best isolate or maximally activate a particular shoulder girdle muscle,^{27,93-96} which exercises elicit coactivation or global activation of many of the muscles that are important for shoulder function,^{30,31,97} and which mode of exercise (resistance tubing, free weights, oscillatory devices, closed kinetic chain)^{29-31,97-103} is most effective in inducing the desired muscle activation. Unfortunately, many of these studies are performed with convenience samples of participants who possess no shoulder injury. Given the demonstrated sensorimotor deficits we see with injury, it is unclear whether these activation patterns reported in healthy study participants can be expected in an injured population as well. However, both Kibler et al⁹³ and Ludewig et al³⁰ demonstrated no scapular muscle activation differences between symptomatic and asymptomatic shoulders during performance of scapular stabilization exercises, suggesting that similar patterns of activation may exist in injured individuals such as the healthy convenience sample participants often assessed. In general, the level of efficacy for using sensorimotor training exercises to elicit desired activation patterns might be improved by demonstrating similar patterns in shoulders with injury, through controlled research studies that use both injured and uninjured populations.

In addition to the measured acute adaptations to the

sensorimotor system, we found 9 orthopedic or sports-related articles that met the inclusion criteria for PEDro and demonstrated the presence or absence of chronic changes to the sensorimotor system resulting from performing sensorimotor training exercises over an extended period of time (Table 2). A majority of those articles were given a PEDro score of 6 (of 10), with a lack of blinding of participants, therapists, or assessors as the most common reason for deduction of the PEDro score. Of the studies that were not included in this review because they did not meet the 5 criteria necessary for PEDro inclusion, the most common flaws were that the study did not compare at least 2 interventions (ie, 1 intervention compared with a control or sham, or 2 interventions) and the trial did not involve some attempt at random allocation.

Shoulder plyometric exercises have been shown in several studies we reviewed to have benefits for the sensorimotor system, including improved proprioception, strength, power, and functional performance.¹⁰⁴⁻¹⁰⁶ Similarly, traditional weight room exercises have been shown to result in improved proprioception accompanying strength gains.¹⁰⁷ Padua et al¹⁰⁸ examined whether training with different modes of shoulder strengthening exercises (open kinetic chain tubing, closed kinetic chain stability, and proprioceptive neuromuscular facilitation [PNF] diagonal manual resistance exercises) influence proprioception, single-arm dynamic stability, shoulder-rotation strength, and functional throwing performance in healthy individuals. They showed that the shoulder rotation strength was improved with only 5 weeks of training healthy individuals using either open kinetic chain tubing or PNF exercises. The fact that changes in strength occurred in such a short time (only 5 weeks) suggests that these changes were a result of sensorimotor adaptation given that strength adaptations due to muscle hypertrophy tend to take longer periods of training (8-12 weeks) to manifest.¹⁰⁹ Of note, the study by Padua et al¹⁰⁸ was not able to demonstrate proprioceptive improvements but did demonstrate throwing function performance with the proprioceptive neuromuscular facilitation exercise. Ubinger et al¹¹⁰ demonstrated similar adaptations from performing a 4-week sensorimotor closed kinetic chain training program for the shoulder, showing improvements in shoulder neuromuscular control (as measured through a single arm stability test). These results suggest the sensorimotor system is trainable in a short period of time (4 to 5 weeks).

Ginn and Cohen¹¹¹ focused on the influence of neuro-

muscular training on pain by comparing the effectiveness of exercise therapy aimed at restoring neuromuscular control mechanisms at the shoulder with other conservative interventions for the treatment of chronic shoulder pain with and without accompanying stiffness. Using a randomized control trial, participants either received exercise therapy aimed at restoring dynamic stabilizing mechanisms and muscle coordination at the shoulder, subacromial corticosteroid injection, or a combination of physical modalities and range of motion exercises. At 5 weeks, all participants' pain intensity, functional impairment, active range of motion, isometric muscle force significantly improved with no difference between the treatment groups. The authors concluded that exercise therapy aimed at restoring neuromuscular control, corticosteroid injection, and multiple physical modalities and range of motion exercises are equally effective in the short-term treatment of shoulder pain. Swanik et al¹¹² used functional training that included elastic tubing, prone dumbbell, and closed kinetic chain exercises to reduce the incidence of shoulder pain in collegiate swimmers. From these results, it appears that sensorimotor training has a positive effect on reducing or preventing shoulder pain.

Although evidence suggests that variables like proprioception, strength, power, pain level, and function can be improved by various forms of sensorimotor training, none of the studies that met the inclusion criteria demonstrated actual changes in muscle activation characteristics (eg, activation patterns, coordination, reflex characteristics) or upper extremity movement patterns. For example, Wang et al¹¹³ reported that scapular movement patterns can be changed with exercise. However, their study was not included in this review because it did not meet the PEDro inclusion criteria due to the lack of a control group. Unfortunately, the current literature base is void of strong intervention studies that specifically demonstrate changes in either muscle activation characteristics or improved movement patterns. Ultimately, the next logical step is to develop and validate sensorimotor training programs that change muscle activation and the resulting movement patterns in patients who would benefit from the training after injury or individuals who are susceptible to sustaining shoulder injury as preventive measures. Demonstration of improvements in sensorimotor function in these population using a well-designed experimental studies would provide higher level of evidence toward the effectiveness of the sensorimotor

training in injury intervention. One area that warrants acknowledgement is that a majority of the trials performed to date involved individuals who are currently healthy. We included these trials in the review because the participants tested may benefit from the preventive intervention, given that they are physically active. However, future studies should focus on improving sensorimotor characteristics in participants with injury given that these individuals most likely have sensorimotor deficits, as demonstrated in the literature.

CONCLUSION

The sensorimotor system contributes to joint stability and function through its control over the muscles that cross the shoulder joint complex. Injury, pain, and fatigue to the shoulder can affect the sensorimotor system both centrally and peripherally due to tissue trauma, pain, and attenuation or stretching of the tissues, resulting in decreased stability and function. Given the important role the sensorimotor system plays in joint stability and function and how the sensorimotor system is affected by the joint injury, training of the sensorimotor system is crucial in treatment following injuries. The current evidence suggests that the sensorimotor system is trainable. As such, clinicians should include sensorimotor training components in their rehabilitation following injury given the deficits that are present with injury and pain or as part of their prevention programs to reduce the effects of fatigue on the sensorimotor system. ■

REFERENCES

1. Myers JB, Lephart SM. The role of the sensorimotor system in the athletic shoulder. *J Athl Train*. 2000;35:351-363.
2. Lephart SM, Riemann BL, Fu F. Introduction to the sensorimotor system. In: Lephart SM, Fu FH, eds. *Proprioception and Neuromuscular Control in Joint Stability*. Champaign, IL: Human Kinetics; 2000: xvii-xxiv.
3. Sherrington C. *The Integrative Action of the Nervous System*. New York, NY: Scribner's Son; 1906.
4. Riemann BL, Lephart SM. The sensorimotor system, part 1: The physiological basis of functional joint stability. *J Athl Train*. 2002;37:71-77.
5. Johansson H, Pedersen J, Bergenheim M, Djupsjobacka M. Peripheral afferents of the knee: their effects on central mechanisms regulating muscle stiffness, joint stability, and proprioception and coordination. In: Lephart SM, Fu FH, eds. *Proprioception and Neuromuscular Control in Joint Stability*. Champaign, IL: Human Kinetics; 2000:5-22.
6. Cain PR, Mutschler TA, Fu FH, Lee SK. Anterior stability of the glenohumeral joint. A dynamic model. *Am J Sports Med*. 1987;15:144-

- 148.
7. Warner JJP. The gross anatomy of the joint surfaces, ligaments, labrum, and capsule. In: Matsen III FA, Fu FH, Hawkins RJ, eds. *The Shoulder: A Balance of Mobility and Stability*. Rosemont, IL: American Academy of Orthopaedic Surgeons; 1993;pages?. **[Query #2: Please provide the page range of the chapter.]**
 8. Harryman D, Sidles J, Harris S, Masten F. Laxity of the normal glenohumeral joint: A quantitative in-vivo assessment. *J Shoulder Elbow Surg*. 1990;1:66-76.
 9. Riemann BL, Lephart SM. The sensorimotor system, part 2: The role of proprioception in motor control and functional joint stability. *J Athl Train*. 2002;37:80-84.
 10. Sjolander P, Johansson H, Djupsjobacka M. Spinal and supraspinal effects of activity in ligament afferents. *J Electromyogr Kinesiol*. 2002;12:167-176.
 11. Myers JB, Riemann BL, Ju YY, Hwang JH, McMahon PJ, Lephart SM. Shoulder muscle reflex latencies under various levels of muscle contraction. *Clin Orthop*. 2003;402:92-101.
 12. Huxel KC, Swanik CB, Swanik KA, et al. Stiffness regulation and muscle-recruitment strategies of the shoulder in response to external rotation perturbations. *J Bone Joint Surg Am*. 2008;90:154-162.
 13. Nyland JA, Caborn DN, Johnson DL. The human glenohumeral joint: a proprioceptive and stability alliance. *Knee Surg Sports Traumatol Arthrosc*. 1998;6:50-61.
 14. Clark JM, Harryman DT II. Tendons, ligaments, and capsule of the rotator cuff. Gross and microscopic anatomy. *J Bone Joint Surg Am*. 1992;74:713-725.
 15. Lee SB, Kim KJ, O'Driscoll SW, Morrey BF, An KN. Dynamic glenohumeral stability provided by the rotator cuff muscles in the mid-range and end-range of motion. A study in cadavera. *J Bone Joint Surg Am*. 2000;82:849-857.
 16. Lippitt S, Matsen F. Mechanisms of glenohumeral joint stability. *Clin Orthop*. 1993;(291):20-28.
 17. Lippitt S, Vanderhoof J, Harris S, Sidles J, Harryman D, Matsen F. Glenohumeral stability from concavity-compression: a quantitative analysis. *J Shoulder Elbow Surg*. 1993;2:27-34.
 18. Halder AM, Halder CG, Zhao KD, O'Driscoll SW, Morrey BF, An KN. Dynamic inferior stabilizers of the shoulder joint. *Clin Biomech*. 2001;16:138-143.
 19. Halder AM, O'Driscoll SW, Heers G, et al. Biomechanical comparisons of effects of supraspinatus tendon detachment, tendon defects, and muscle retraction. *J Bone Joint Surg Am*. 2002;84:780-785.
 20. Halder AM, Zhao KD, Odriscoll SW, Morrey BF, An KN. Dynamic contributions to superior shoulder stability. *J Orthop Res*. 2001;19:206-212.
 21. Parsons IM, Apreleva M, Fu FH, Woo SLY. The effect of rotator cuff tears on reaction forces at the glenohumeral joint. *J Orthop Res*. 2002;20:439-446.
 22. Labriola JE, Lee TQ, Debski RE, McMahon PJ. Stability and instability of the glenohumeral joint: the role of shoulder muscles. *J Shoulder Elbow Surg*. 2005;14(suppl):325-385.
 23. Wuelker N, Korell M, Thren K. Dynamic glenohumeral joint stability. *J Shoulder Elbow Surg*. 1998;7:43-52.
 24. Sharkey NA, Marder RA. The rotator cuff opposes superior translation of the humeral head. *Am J Sports Med*. 1995;23:270-275.
 25. Myers JB, Hwang JH, Pasquale MR, Blackburn JT, Lephart SM. Rotator cuff coactivation ratios in participants with subacromial impingement syndrome [published online ahead of print August 31, 2008]. *J Sci Med Sport*. **[Query #3: Please provide the doi, if possible.]**
 26. Moseley JB, Jobe FW, Pink M, Perry J, Tibone J. EMG analysis of the scapular muscles during a shoulder rehabilitation program. *Am J Sports Med*. 1992;20:128-134.
 27. Decker MJ, Hintermeister RA, Faber KJ, Hawkins RJ. Serratus anterior muscle activity during selected rehabilitation exercises. *Am J Sports Med*. 1999;27:784-791.
 28. Ekstrom RA, Bifulco KM, Lopau CJ, Andersen CF, Gough JR. Comparing the function of the upper and lower parts of the serratus anterior muscle using surface electromyography. *J Orthop Sports Phys Ther*. 2004;34:235-243.
 29. Ekstrom RA, Donatelli RA, Soderberg GL. Surface electromyographic analysis of exercises for the trapezius and serratus anterior muscles. *J Orthop Sports Phys Ther*. 2003;33:247-258.
 30. Ludewig PM, Hoff MS, Osowski EE, Meschke SA, Rundquist PJ. Relative balance of serratus anterior and upper trapezius muscle activity during push-up exercises. *Am J Sports Med*. 2004;32:484-493.
 31. Myers JB, Pasquale MR, Laudner KG, Sell TC, Bradley JP, Lephart SM. On-the-field resistance tubing exercises for throwers: An electromyographic analysis. *J Athl Train*. 2005;40:15-22.
 32. Kibler WB. The role of the scapula in athletic shoulder function. *Am J Sports Med*. 1998;26:325-337.
 33. Ervilha UF, Farina D, Arendt-Nielsen L, Graven-Nielsen T. Experimental muscle pain changes motor control strategies in dynamic contractions. *Exp Brain Res*. 2005;164:215-224.
 34. Falla D, Farina D, Graven-Nielsen T. Experimental muscle pain results in reorganization of coordination among trapezius muscle subdivisions during repetitive shoulder flexion. *Exp Brain Res*. 2007;178:385-393.
 35. Farina D, Arendt-Nielsen L, Merletti R, Graven-Nielsen T. Effect of experimental muscle pain on motor unit firing rate and conduction velocity. *J Neurophysiol*. 2004;91:1250-1259.
 36. Graven-Nielsen T, Lund H, Arendt-Nielsen L, Danneskiold-Samsøe B, Bliddal H. Inhibition of maximal voluntary contraction force by experimental muscle pain: A centrally mediated mechanism. *Muscle & Nerve*. 2002;26:708-712.
 37. Graven-Nielsen T, Svensson P, Arendt-Nielsen L. Effects of experimental muscle pain on muscle activity and co-ordination during static and dynamic motor function. *Electroencephalogr Clin Neurophysiol* 1997;105:156-164.
 38. Le Pera D, Graven-Nielsen T, Valeriani M, et al. Inhibition of motor system excitability at cortical and spinal level by tonic muscle pain. *Clin Neurophysiol*. 2001;112:1633-1641.
 39. Hess SA, Richardson C, Darnell R, Fries P, Lisle D, Myers P. Timing of rotator cuff activation during shoulder external rotation in throwers with and without symptoms of pain. *J Orthop Sports Phys Ther*. 2005;35:812-820.
 40. Steenbrink F, de Groot JH, Veeger HE, Meskers CG, van de Sande MA, Rozing PM. Pathological muscle activation patterns in patients with massive rotator cuff tears, with and without subacromial anaesthetics. *Manual Ther*. 2006;11:231-237.
 41. Scibek JS, Mell AG, Downie BK, Carpenter JE, Hughes RE. Shoulder kinematics in patients with full-thickness rotator cuff tears after a subacromial injection. *J Shoulder Elbow Surg*. 2008;17:172-181.
 42. Lephart S, Warner J, Borsa P, Fu F. Proprioception of the shoulder joint in healthy, unstable, and surgically repaired shoulders. *J*

- Shoulder Elbow Surg.* 1994;3:371-380.
43. Smith RL, Brunolli J. Shoulder kinesthesia after anterior glenohumeral dislocation. *Phys Ther.* 1989;69:106-112.
 44. Zuckerman JD, Gallagher MA, Cuomo F, Rokito A. The effect of instability and subsequent anterior shoulder repair on proprioceptive ability. *J Shoulder Elbow Surg.* 2003;12:105-109.
 45. Barden JM, Balyk R, Raso VJ, Moreau M, Bagnall K. Atypical shoulder muscle activation in multidirectional instability. *Clin Neurophysiol.* 2005;116:1846-1857.
 46. Cools AM, Declercq GA, Cambier DC, Mahieu NN, Witvrouw EE. Trapezius activity and intramuscular balance during isokinetic exercise in overhead athletes with impingement symptoms. *Scand J Med Sci Sports.* 2007;17:25-33.
 47. de Moraes Faria CD, Teixeira-Salmela LF, de Paula Goulart FR, de Souza Moraes GF. Scapular muscular activity with shoulder impingement syndrome during lowering of the arms. *Clin J Sport Med.* 2008;18:130-136.
 48. Hundza SR, Zehr EP. Muscle activation and cutaneous reflex modulation during rhythmic and discrete arm tasks in orthopaedic shoulder instability. *Exp Brain Res.* 2007;179:339-351.
 49. Myers JB, Ju YY, Hwang JH, McMahon PJ, Rodosky MW, Lephart SM. Reflexive muscle activation alterations in shoulders with anterior glenohumeral instability. *Am J Sports Med.* 2004;32:1013-1021.
 50. Matias R, Pascoal AG. The unstable shoulder in arm elevation: A three-dimensional and electromyographic study in subjects with glenohumeral instability. *Clin Biomech.* 2006;21(suppl 1):S52-S58.
 51. Lephart SM, Pincivero DM, Giraldo JL, Fu FH. The role of proprioception in the management and rehabilitation of athletic injuries. *Am J Sports Med.* 1997;25:130-137.
 52. Tibone JE, Fechter J, Kao JT. Evaluation of a proprioception pathway in patients with stable and unstable shoulders with somatosensory cortical evoked potentials. *J Shoulder Elbow Surg.* 1997;6:440-443.
 53. Barden JM, Balyk R, Raso VJ, Moreau M, Bagnall K. Dynamic upper limb proprioception in multidirectional shoulder instability. *Clin Orthop.* 2004;(420):181-189.
 54. Illyes A, Kiss RM. Kinematic and muscle activity characteristics of multidirectional shoulder joint instability during elevation. *Knee Surg Sports Traumatol Arthrosc.* 2006;14:673-685.
 55. Morris AD, Kemp GJ, Frostick SP. Shoulder electromyography in multidirectional instability. *J Shoulder Elbow Surg.* 2004;13:24-29.
 56. Allegrucci M, Whitney SL, Lephart SM, Irrgang JJ, Fu FH. Shoulder kinesthesia in healthy unilateral athletes participating in upper extremity sports. *J Orthop Sports Phys Ther.* 1995;21:220-226.
 57. Safran MR, Borsa PA, Lephart SM, Fu FH, Warner JJ. Shoulder proprioception in baseball pitchers. *J Shoulder Elbow Surg.* 2001;10:438-444.
 58. Blasier RB, Carpenter JE, Huston LJ. Shoulder proprioception: Effects of joint laxity, joint position, and direction of motion. *Orthop Rev.* 1994;23:45-50.
 59. Lephart SM, Myers JB, Bradley JP, Fu FH. Shoulder proprioception and function following thermal capsulorrhaphy. *Arthroscopy.* 2002;18:770-778.
 60. Sullivan JA, Hoffman MA, Harter RA. Shoulder joint position sense after thermal, open, and arthroscopic capsulorrhaphy for recurrent anterior instability. *J Shoulder Elbow Surg.* 2008;17:389-394.
 61. Bandholm T, Rasmussen L, Aagaard P, Jensen BR, Diederichsen L. Force steadiness, muscle activity, and maximal muscle strength in subjects with subacromial impingement syndrome. *Muscle & Nerve.* 2006;34:631-639.
 62. Clisby EF, Bitter NL, Sandow MJ, Jones MA, Magarey ME, Jaberzadeh S. Relative contributions of the infraspinatus and deltoid during external rotation in patients with symptomatic subacromial impingement. *J Shoulder Elbow Surg.* 2008;17(suppl):87S-92S.
 63. Kido T, Itoi E, Konno N, Sano A, Urayama M, Sato K. Electromyographic activities of the biceps during arm elevation in shoulders with rotator cuff tears. *Acta Orthop Scand.* 1998;69:575-579.
 64. Kido T, Itoi E, Konno N, Sano A, Urayama M, Sato K. The depressor function of biceps on the head of the humerus in shoulders with tears of the rotator cuff. *J Bone Joint Surg Br.* 2000;82:416-419.
 65. Kelly BT, Williams RJ, Cordasco FA, et al. Differential patterns of muscle activation in patients with symptomatic and asymptomatic rotator cuff tears. *J Shoulder Elbow Surg.* 2005;14:165-171.
 66. Moraes GF, Faria CD, Teixeira-Salmela LF. Scapular muscle recruitment patterns and isokinetic strength ratios of the shoulder rotator muscles in individuals with and without impingement syndrome. *J Shoulder Elbow Surg.* 2008;17(suppl):48S-53S.
 67. Voight ML, Hardin JA, Blackburn TA, Tippet S, Canner GC. The effects of muscle fatigue on and the relationship of arm dominance to shoulder proprioception. *J Orthop Sport Phys Ther.* 1996;23:348-352.
 68. Myers JB, Guskiewicz KM, Schneider RA, Prentice WE. Proprioception and neuromuscular control of the shoulder after muscle fatigue. *J Athl Training.* 1999;34:362-367.
 69. Tripp BL, Boswell L, Gansnedder BM, Shultz SJ. Functional fatigue decreases 3-dimensional multijoint position reproduction acuity in the overhead-throwing athlete. *J Athl Train.* 2004;39:316-320.
 70. Tripp BL, Uhl TL, Mattacola CG, Srinivasan C, Shapiro R. Functional multijoint position reproduction acuity in overhead-throwing athletes. *J Athl Train.* 2006;41:146-153.
 71. Tripp BL, Yochem EM, Uhl TL. Recovery of upper extremity sensorimotor system acuity in baseball athletes after a throwing-fatigue protocol. *J Athl Train.* 2007;42:452-457.
 72. Tripp BL, Yochem EM, Uhl TL. Functional fatigue and upper extremity sensorimotor system acuity in baseball athletes. *J Athl Train.* 2007;42:90-98.
 73. Lee HM, Liao JJ, Cheng CK, Tan CM, Shih JT. Evaluation of shoulder proprioception following muscle fatigue. *Clin Biomech.* 2003;18:843-847.
 74. Carpenter JE, Blasier RB, Pellizzon GG. The effects of muscle fatigue on shoulder joint position sense. *Am J Sports Med.* 1998;26:262-264.
 75. Pedersen J, Lonn J, Hellstrom F, Djupsjobacka M, Johansson H. Localized muscle fatigue decreases the acuity of the movement sense in the human shoulder. *Med Sci Sports Exerc.* 1999;31:1047-1052.
 76. McQuade KJ, Hwa Wei S, Smidt GL. Effects of local muscle fatigue on three-dimensional scapulohumeral rhythm. *Clin Biomech.* 1995;10:144-148.
 77. Bowman TG, Hart JM, McGuire BA, Palmieri RM, Ingersoll CD. A functional fatiguing protocol and deceleration time of the shoulder from an internal rotation perturbation. *J Athl Train.* 2006;41:275-279.
 78. Cools AM, Witvrouw EE, De Clercq GA, et al. Scapular muscle recruitment pattern: Electromyographic response of the trapezius muscle to sudden shoulder movement before and after a fatiguing exercise. *J Orthop Sports Phys Ther.* 2002;32:221-229.
 79. Minning S, Eliot CA, Uhl TL, Malone TR. EMG analysis of shoulder

- muscle fatigue during resisted isometric shoulder elevation. *J Electromyogr Kinesiol*. 2007;17:153-159.
80. Rodrigues D, Pessan MA, Kawano MM, Stabile GR, Cardoso JR. Electromyographic analysis of deltoid muscle fatigue during abduction on scapular and frontal planes. *Electromyogr Clin Neurophysiol*. 2008;48:293-300.
 81. Kawczynski A, Nie H, Jaskolska A, Jaskolski A, Arendt-Nielsen L, Madeleine P. Mechanomyography and electromyography during and after fatiguing shoulder eccentric contractions in males and females. *Scand J Med Sci Sports*. 2007;17:172-179.
 82. Rodgers MM, Gayle GW, Figoni SF, Kobayashi M, Lieh J, Glaser RM. Biomechanics of wheelchair propulsion during fatigue. *Arch Phys Med Rehabil*. 1994;75:85-93.
 83. Rodgers MM, McQuade KJ, Rasch EK, Keyser RE, Finley MA. Upper-limb fatigue-related joint power shifts in experienced wheelchair users and nonwheelchair users. *J Rehabil Res Dev*. 2003;40:27-37.
 84. Cote JN, Feldman AG, Mathieu PA, Levin MF. Effects of fatigue on intermuscular coordination during repetitive hammering. *Motor Control*. 2008;12:79-92.
 85. Cote JN, Raymond D, Mathieu PA, Feldman AG, Levin MF. Differences in multi-joint kinematic patterns of repetitive hammering in healthy, fatigued and shoulder-injured individuals. *Clin Biomech*. 2005;20:581-590.
 86. Teyhen DS, Miller JM, Middag TR, Kane EJ. Rotator cuff fatigue and glenohumeral kinematics in participants without shoulder dysfunction. *J Athl Train*. 2008;43:352-358.
 87. Murray TA CT, Werner SL, Schlegel TF, Hawkins RJ. The effects of extended play on professional baseball pitchers. *Am J Sports Med*. 2001;29:137-142.
 88. McQuade KJ, Dawson J, Smidt GL. Scapulothoracic muscle fatigue associated with alterations in scapulohumeral rhythm kinematics during maximum resistive shoulder elevation. *J Orthop Sports Phys Ther*. 1998;28:74-80.
 89. Tsai NT, McClure PW, Karduna AR. Effects of muscle fatigue on 3-dimensional scapular kinematics. *Arch Phys Med Rehabil*. 2003;84:1000-1005.
 90. Su KP, Johnson MP, Gracely EJ, Karduna AR. Scapular rotation in swimmers with and without impingement syndrome: Practice effects. *Med Sci Sports Exerc*. 2004;36:1117-1123.
 91. Ebaugh DD, McClure PW, Karduna AR. Effects of shoulder muscle fatigue caused by repetitive overhead activities on scapulothoracic and glenohumeral kinematics. *J Electromyogr Kinesiol*. 2006;16:224-235.
 92. Ebaugh DD, McClure PW, Karduna AR. Scapulothoracic and glenohumeral kinematics following an external rotation fatigue protocol. *J Orthop Sports Phys Ther*. 2006;36:557-571.
 93. Kibler WB, Sciascia AD, Uhl TL, Tambay N, Cunningham T. Electromyographic analysis of specific exercises for scapular control in early phases of shoulder rehabilitation. *Am J Sports Med*. 2008;36:1789-1798.
 94. Decker MJ, Tokish JM, Ellis HB, Torry MR, Hawkins RJ. Subscapularis muscle activity during selected rehabilitation exercises. *Am J Sports Med*. 2003;31:126-134.
 95. Hintermeister RA, Lange GW, Schultheis JM, Bey MJ, Hawkins RJ. Electromyographic activity and applied load during shoulder rehabilitation exercises using elastic resistance. *Am J Sports Med*. 1998;26:210-220.
 96. Reddy AS, Mohr KJ, Pink MM, Jobe FW. Electromyographic analysis of the deltoid and rotator cuff muscles in persons with subacromial impingement. *J Shoulder Elbow Surg*. 2000;9:519-523.
 97. Uhl TL, Carver TJ, Mattacola CG, Mair SD, Nitz AJ. Shoulder musculature activation during upper extremity weight-bearing exercise. *J Orthop Sports Phys Ther*. 2003;33:109-117.
 98. Lehman GJ, Gilas D, Patel U. An unstable support surface does not increase scapulothoracic stabilizing muscle activity during push up and push up plus exercises. *Manual Ther*. 2008;13:500-506.
 99. Lehman GJ, MacMillan B, MacIntyre I, Chivers M, Fluter M. Shoulder muscle EMG activity during push up variations on and off a Swiss ball. *Dyn Med*. 2006;5:7.
 100. Lister JL, Del Rossi G, Ma F, et al. Scapular stabilization activity during body blade, cuff weights, and thera-band use. *J Sport Rehabil*. 2007;16:50-67.
 101. Kinney E, Wusthoff J, Zyck A, et al. Activation of the trapezius muscle during varied forms of Kendall exercises. *Phys Ther Sport*. 2008;9:3-8.
 102. Reinold MM, Macrina LC, Wilk KE, et al. Electromyographic analysis of the supraspinatus and deltoid muscles during 3 common rehabilitation exercises. *J Athl Train*. 2007;42:464-469.
 103. Reinold MM, Wilk KE, Fleisig GS, et al. Electromyographic analysis of the rotator cuff and deltoid musculature during common shoulder external rotation exercises. *J Orthop Sports Phys Ther*. 2004;34:385-394.
 104. Carter AB, Kaminski TW, Douex AT, Jr, Knight CA, Richards JG. Effects of high volume upper extremity plyometric training on throwing velocity and functional strength ratios of the shoulder rotators in collegiate baseball players. *J Strength Cond Res*. 2007;21:208-215.
 105. Schulte-Edelmann JA, Davies GJ, Kernozek TW, Gerberding ED. The effects of plyometric training of the posterior shoulder and elbow. *J Strength Cond Res*. 2005;19:129-134.
 106. Swanik KA, Lephart SM, Swanik CB, Lephart SP, Stone DA, Fu FH. The effects of shoulder plyometric training on proprioception and selected muscle performance characteristics. *J Shoulder Elbow Surg*. 2002;11:579-586.
 107. Rogol IM, Ernst GP, Perrin DH. Open and closed kinetic chain exercises improve shoulder joint reposition sense equally in healthy subjects. *J Athl Train*. 1998;33:315-318.
 108. Padua DA, Guskiewicz KM, Prentice WE, Schneider RA, Shields EW. The effect of select shoulder exercises on strength, active angle reproduction, single-arm balance, and functional performance. *J Sport Rehabil*. 2004;13:75-95.
 109. Moritani T, deVries HA. Neural factors versus hypertrophy in the time course of muscle strength gain. *Am J Phys Med*. 1979;58:115-130.
 110. Ubinger ME, Prentice WE, Guskiewicz KM. Effects of closed kinetic chain training on neuromuscular control in the upper extremity. *J Sport Rehabil*. 1999;8:184-194.
 111. Ginn KA, Cohen ML. Exercise therapy for shoulder pain aimed at restoring neuromuscular control: A randomized comparative clinical trial. *J Rehabil Med*. 2005;37:115-122.
 112. Swanik KA, Swanik CB, Lephart SM, Huxel K. The effect of functional training on the incidence of shoulder pain and strength in intercollegiate swimmers. *J Sport Rehabil*. 2002;11:140-154.
 113. Wang CH, McClure P, Pratt NE, Nobiletti R. Stretching and strengthening exercises: Their effect on three-dimensional scapular kinematics. *Arch Phys Med Rehabil*. 1999;80:923-929.
 114. Kraemer WJ, Ratamess N, Fry AC, et al. Influence of resistance training volume and periodization on physiological and performance adaptations in collegiate women tennis players. *Am J Sports Med*.