

# A New Paradigm: A Joint Test of Structural and Correlation Parameters in Instrumental Variables Regression When Perfect Exogeneity is Violated

*By* MEHMET CANER AND MELINDA SANDLER MORRILL \*

September 22, 2009

## Abstract

Currently, the commonly employed instrumental variables strategy relies on the knife-edge assumption of perfect exogeneity for valid inference. To make reliable inferences on the structural parameters under violations of exogeneity one must know the true correlation between the structural error and the instruments. The main innovation in this paper is to identify an appropriate test in this context: a joint null hypothesis of the structural parameters with the correlation between the instruments and the structural error term. We introduce a new endogeneity accounted test by combining the structural parameter inference while correcting the bias associated with nonexogeneity of instrument. To address inference under violations of exogeneity, significant contributions have been made in the recent literature by assuming some degree of non-exogeneity. A key advantage of our approach over that of the previous literature is that we do not need to make any assumptions about the degree of violation of exogeneity either as possible values or prior distributions. In particular, our method is not a form of sensitivity analysis. Since our test statistic is continuous and monotonic in correlation one can conduct inference for structural parameters by a simple grid search over correlation values. We can make accurate inferences on the structural parameters because of a feature of the grid search over correlation values. One can also build joint confidence intervals for the structural parameters and the correlation parameter by inverting the test statistic. In the inversion, the null values of these parameters are used. We also propose a new way of testing exclusion restrictions, even in the just identified case.

## 1 Introduction

Economists frequently apply instrumental variable methods to draw inferences about whether or not a variable influences an economic outcome. For example, labor economists employ varied

---

\*Caner: North Carolina State University, Department of Economics, 4168 Nelson Hall, Raleigh, NC 27518 (email: mcaner@ncsu.edu). Morrill: North Carolina State University, Department of Economics, 4144 Nelson Hall, Raleigh, NC 27695-8110 (email: melinda\_morrill@ncsu.edu). We thank Daron Acemoglu for his comments and encouragement. We thank Josh Angrist and Atsushi Inoue for insightful comments and suggestions. We also thank Tengying Weng who provided research assistance.

instruments, including quarter and year of birth (Angrist and Krueger, 1991), tuition and distance to nearest college (Kane and Rouse, 1995, Card, 1993), attending reform school (Meghir and Palene, 2005) and birth year interacted with school buildings in region of birth (Dufflo, 2001) to measure the extent to which a person's education influences her salary and wages. In a distinct but related literature that combines macro-economics, political economy and comparative institutions, economists employ instruments including early settler mortality (Acemoglu, Johnson and Robinson, 2001), ethnic capital (Hall and Jones, 1999), ethno-linguistic fractionalization (Mauro, 1995) and legal families (Djankov et al., 2003, and Acemoglu and Johnson, 2006) to determine whether or not the quality of institutions influences long term growth and investment.

Instrumental variable (IV) methods are used to identify causal relationships by isolating changes in an endogenous variable (or variables) that are unrelated to potential unobserved factors. To identify a causal relationship, instruments must be exogenous; that is, they are not related to the outcome variable after controlling for relevant explanatory variables. For example, early settler mortality is exogenous if it is only related to long term growth through its impact on institutions, after controlling for relevant variables such as latitude. This requirement is strong because it means that settler mortality can only influence long term growth indirectly through the quality of contemporary institutions. The exogeneity of early settler mortality, however, is controversial; for example, as noted by Glaeser et al. (2004), early settler mortality could also influence long term growth through its impact on the unobservable human capital of the early settlers. Whether or not the exclusion restriction is perfectly satisfied is debatable for many (and perhaps most) applications of instrumental variables.

Applied econometricians use a simple t-ratio test statistic to infer whether there is causality among the variables analyzed using IV methods. However, the t-test gives unreliable results even when there is a slight violation of exogeneity, as established recently in a paper by Berkowitz, Caner and Fang (2008). As shown in Table 1, the standard t-ratio has undesirable properties when the instrument is endogenous. The table shows the actual size of the test when the single instrument is endogenous. Larger sample sizes exacerbate the problem. For example, with the true correlation of 0.1 at a sample size of  $n = 100$  the size of the test is 19.9%, while if the sample size increases to  $n = 1000$ , the size is 88.8%. This is a massive size distortion.

In this paper, we propose a new test that modifies the t-test in a very simple way, yet is robust to instrument validity concerns. The main innovation is the recognition that the

appropriate test is actually a joint test of the structural parameters with the correlation between the structural error and the instruments. Since inferences on the structural parameters depend on this correlation, a joint null is needed. The idea of the test is to subtract the drift from the standard t statistic. The drift depends on the value of the true correlation between the structural error and the instrument. Since the true correlation is between -1 and 1 we can do a grid search over all possible values of the correlation.

In one sense, we propose a test that combines structural parameter inference with a test of violations of exogeneity. The test statistic here is useful for the just identified case as well as the over-identified case. Furthermore, our test can be inverted to get valid confidence intervals for the structural parameters ( $\beta_0$ ) and the correlation parameter ( $\rho_0$ ) by doing a joint grid search. We record the values of the test where the null is not rejected as being in the joint confidence interval. We should remind readers that we are not conditioning on one parameter to get the other's confidence interval. The joint confidence interval can be unbounded with positive probability, so this is a valid confidence interval. The test statistic we derive has standard normal distribution when we have only one endogenous regressor. It has  $\chi^2$  limit when the number of endogenous regressors are larger than one. So the test is asymptotically pivotal regardless of exogeneity violations.

One of the key issues is the joint nature of the test. One may be concerned that this would impede inference on the structural parameters alone. In other words, the concern is that one could reject the true null due to usage of the wrong correlation values. In fact, this is not a problem in our context. Since the test statistic is continuous and monotonic in correlation, we will eventually find the true correlation and can conduct inference at that point. If the joint null is true, then the test statistic value will be below the critical values at the true correlation. In that case we do not reject the null and have correct inference. If the alternative is true, our test rejects the false null regardless of correlation values, as does the standard t-test. Therefore even though the joint null may seem to be limiting in interpreting rejection, we can still make accurate inferences on the structural parameters.

We also introduce a new exclusion restriction test. This test is valid even when there are only the same number of instruments as endogenous regressors. The joint null in that case is no correlation and structural parameters taking certain values. By testing several possible grid values of the structural parameter, we have several test statistics recorded. If all of tests reject the null of no correlation with the structural parameter null value that is tested, then the

instrument cannot be exogenous. Simulations show that this has good finite sample properties and can be used in the applied work.

We present two empirical examples that demonstrate the methods we propose. In the first, Acemoglu and Johnson (2005) measure the effects of institutions on country-level gross domestic product (GDP) using settler mortality levels as an instrument for the development of institutions. The exclusion restriction requires that early settler mortality rates only affect GDP through the development of institutions. According to the empirical exercise described in Section 5, we find a region of non-rejection that is clearly bounded away from the structural parameter equaling zero. This is a very powerful result because it demonstrates that, we can still infer that the coefficient on institutions is positive and statistically significant. Since a perfectly exogenous instrument is very difficult (or perhaps impossible) to find, our test allows researchers to understand the regions where the joint null fails to reject, yielding a joint confidence interval for the correlation and structural parameters.

In addition to the Acemoglu and Johnson example, we demonstrate this method using a well-known example from labor economics. In Card (1995), the author argues that proximity to college can be used as an instrument for college attendance when calculating the returns to schooling on wages. One potential violation of exogeneity is that proximity to college is correlated with other unobserved factors that are positively associated with high earnings, such as having well-educated parents or having a higher quality public primary and secondary education. Although the original estimates indicated statistical significance, we demonstrate that in this case the confidence interval for the structural parameter is unbounded in both directions. Section 5 includes a detailed discussion of the interpretation and implication of these results.

Previous to the literature cited below, research using the instrumental variables strategy has been conducted on the unreasonable assumption that perfect exogeneity is satisfied. Recent work in applied econometrics has made large strides by questioning the validity of the perfect exogeneity assumption and demonstrating the serious implications of violations of this assumption. Berkowitz, Caner and Fang (2009) analyze the Anderson-Rubin test with a new resampling scheme when there is violation of exogeneity. The main assumption in that paper is a violation of exogeneity, but this is local to zero. In large samples perfect exogeneity is assumed. In finite samples, the block size choice is important in their resampling scheme.

Other related and important papers include Nevo and Rosen (2009), where they provide analytical bounds on structural parameters via set identification. Ashley (2009) provides a

sensitivity analysis for the instrumental variable regression based on the covariance between the instruments and the structural error. That study shows clearly that correlation plays an important role in inference for structural parameters. He modifies the structural equation so that the new error is orthogonal to the existing instrument. Then he derives the limit for this new two stage least squares estimator. A very recent paper by Guggenberger (2009) considers the several variants of subsampling to gather information about local violation of exogeneity in identification robust tests. He shows that even though the Anderson-Rubin test is oversized, compared to other identification robust test the Anderson-Rubin test performs well. Caner (2009) also considers a similar problem, and shows that Anderson-Rubin test is robust to local violations of exogeneity in the many moments framework. In this paper we do not analyze the Anderson-Rubin test. Rather, we focus on the standard t-test, upon which almost all of the applied literature relies. In the future, we plan to extend our analysis to incorporate modifications to the Anderson-Rubin test.

For Bayesian analysis on the issue of violations of exogeneity in instrumental variables, two good sources are by Conley, Hansen, Rossi (2007) and Kray (2008). Conley, Hansen, and Rossi (2007) consider a three way attack on the problem of violation of exogeneity. The first attempt is to assume a support for the parameter that controls the violation and then use that assumption to build confidence intervals for the structural parameters. This proves to be conservative. The second approach they take is to put a prior on the parameter for the violation and then build confidence intervals for the structural parameters. The third line of attack is fully Bayesian and puts a prior on parameter controlling the violation of exogeneity but allows this parameter to be independent from the structural parameters priors. Then a variant of that is to assume the priors are dependent on each other.

The method presented in this paper take an entirely new approach to the problem of non-exogeneity in instrumental variables estimation. To our best knowledge, our paper is the first to analyze the correlation parameter and the structural parameters jointly. The previous literature conditioned on the correlation parameter in one way and then conducted inference on structural parameters. That approach is valuable, but it assumes we have some knowledge of the degree of violation of exogeneity. In this paper we build a joint test for structural parameters and the correlation parameter. This changes the paradigm of inference in instrumental variable regression and provides a framework that is tractable and straightforward to implement empirically.

The remaining sections of this paper are as follows. Section 2 provides a theoretical basis for

the empirical technique. In Section 3, we present an algorithm that suggests a new exogeneity accounted test that reflects the joint nature of the null hypothesis. Section 4 presents simulation results which justify the practicality and efficiency of our estimates. Section 5 presents the two examples of the application of this technique in empirical research. Section 6 provides a discussion and concludes. The proofs are included in Appendix A, and the Stata code used to implement the test statistic is included in Appendix B.

## 2 The Model and Assumptions

We consider the following linear simultaneous equations model:

$$y = X\beta_0 + u, \tag{1}$$

$$X = Z\pi_0 + V, \tag{2}$$

where we refer to (1) as the structural equation and (2) as the first stage equation. Here,  $X$  represents  $n \times k$  matrix,  $X_i$  is a  $k \times 1$  vector of endogenous variables, and  $Z$  is  $n \times l$  matrix. Below,  $Z_i$  is  $l \times 1$  vector of instruments,  $l \geq k$ . Also  $\pi_0$  is of full column rank  $k$ . The errors  $u_i, V_{ij}, i = 1, \dots, n, j = 1, \dots, k$  are correlated. Control variables may be added to the system. If this is the case, one can simply project them out and the analysis below follows. The variance matrix  $EV_iV_i' = \Sigma_{VV} < \infty$ , and nonsingular.  $Eu_i^2 = \sigma_u^2 < \infty$ , and  $Eu_i = EV_i = 0$ . Let  $\hat{\beta}$  represent the two-stage least squares (2SLS) estimate of  $\beta_0$ , and  $\hat{\pi}$  is the least squares (LS) estimate of  $\pi_0$ .

**Assumption 1.** (i). (Violation of Exogeneity)

$$EZ_iu_i = C,$$

where  $C$  is  $l \times 1$  vector with  $C = (C_1, \dots, C_m, \dots, C_l)'$ , and each  $C_m \neq 0$  and is finite for  $m = 1, 2, \dots, l$ .

(ii). We also have

$$EZ_iV_i' = 0.$$

(iii).  $EZ_iX_i'$  has full column rank.

**Assumption 2.** The following limits hold jointly when the sample size  $n$  converges to infinity:

(i).

$$\left( n^{-1} \sum_{i=1}^n u_i^2, n^{-1} \sum_{i=1}^n V_i u_i, n^{-1} \sum_{i=1}^n V_i V_i' \right) \xrightarrow{p} (\sigma_u^2, \Sigma_{Vu}, \Sigma_{VV}),$$

where  $\sigma_u^2, \Sigma_{Vu}, \Sigma_{VV}$  are scalar,  $k \times 1$ , and  $k \times k$ , respectively. The scalar is positive, the vector is nonzero and finite, and the matrix is positive definite and finite.

(ii). We have the following law of large numbers

$$\hat{Q}_{zz} = n^{-1} \sum_{i=1}^n Z_i Z_i' \xrightarrow{p} Q_{zz},$$

where  $Q_{zz}$  is a positive definite and finite  $k \times k$  matrix.

(iii). We have the following central limit theorem

$$\left( n^{-1/2} \sum_{i=1}^n (Z_i u_i - E Z_i u_i), n^{-1/2} \sum_{i=1}^n Z_i V_i' \right) \xrightarrow{d} (\Psi_{zu}, \Psi_{zV}),$$

$$\begin{pmatrix} \Psi_{zu} \\ \Psi_{zV} \end{pmatrix} \equiv N[0, \Sigma \otimes Q_{zz}],$$

and

$$\Sigma = \begin{pmatrix} \sigma_u^2 & \Sigma'_{Vu} \\ \Sigma_{Vu} & \Sigma_{VV} \end{pmatrix}.$$

Note that Assumption 1i is the main issue of this paper. The perfect exogeneity that is used in instrumental variable analysis is a knife-edge, unrealistic assumption for applied work. Even though the researcher is careful in selecting the “perfectly exogenous” instrument there can still be unavoidable violations of exogeneity. There will be more discussion about that assumption in the next section.

Another possibility is the case of near exogeneity (a local to zero violation). This is analyzed in Berkowitz, Caner and Fang (2008) and it is shown that the t-test is affected and inference becomes unreliable. But there is no solution that is proposed in that paper. In this paper, with a more realistic assumption, we propose a solution to inference under Assumption 1.

Assumption 2 is basically law of large numbers and a central limit theorem. These hold under primitive conditions, such as moment conditions on the instruments and the errors, for these see Davidson (1994).

### 3 Test Statistics

In this section we discuss and analyze Assumption 1i and based on that introduce three cases of interest in applied work. First we cover the most widely used case of a just identified system with one endogenous regressor and one instrument ( $k = l = 1$ ). Then in our second case we consider one endogenous variable with more than one instrument ( $k = 1, l \geq k$ ). The last case involves the general case where we may have more than one endogenous variable and more than one instrument ( $k > 1, l \geq k$ ).

#### 3.1 The Just Identified Case with One Endogenous Regressor

Since we do not know the true correlation between the instrument and the structural error, and this affects the inference for the structural parameters, we need a joint test of the structural parameter and the correlation between the structural error (second stage error) and the instrument ( $Z_i, u_i$ ). The joint null is  $H_0 : \beta = \beta_0, \rho = \rho_0$ , where  $\rho$  represents the correlation between the structural error and the instrument.

We start with two issues that are related to the standard t statistic. First, the unobserved covariance between the instruments and the structural error can be converted to a measure involving correlation. Because the correlation is standardized so that it is bounded by -1 and +1, we can conduct a finite grid search over potential correlation values. In that respect, assume  $cov(Z_i, u_i) = C$  for all  $i = 1, \dots, n$ , where  $C$  is scalar since  $k = l = 1$ .

Notice that:

$$corr(Z_i, u_i) = \frac{cov(Z_i, u_i)}{\sigma_u \sqrt{var(Z_i)}}.$$

Then,

$$C = \sigma_u \sigma_{zz} \rho_0, \tag{3}$$

where  $\rho_0$  denotes the true unknown correlation, and  $var Z_i = \sigma_{zz}^2$  for all  $i = 1, \dots, n$ .

The second issue is that in the regular t-test, the estimator  $\hat{\sigma}_u^2$  is inconsistent, which is shown in (24) below. When we impose from the null  $\beta = \beta_0$ , we have the following consistent estimate:

$$\tilde{\sigma}_u^2 = n^{-1} \sum_{i=1}^n (y_i - X_i \beta_0)^2 \xrightarrow{p} E u_i^2 = \sigma_u^2.$$

We propose a test statistic evaluated at true correlation ( $\rho_0$ ) and structural parameter ( $\beta_0$ ), denoted as  $NT(\beta_0, \rho_0)$ :

$$NT(\beta_0, \rho_0) = \frac{\sqrt{n}(\hat{\beta} - \beta_0)}{\tilde{\sigma}_u[|\hat{\pi}|^{-1}\hat{Q}_{zz}^{-1/2}]} - \sqrt{n}sgn(\hat{\pi})f(z)\rho_0, \quad (4)$$

where  $sgn(\hat{\pi})$  is the sign of the least squares estimate  $\hat{\pi}$  in (2) for the scalar case, and

$$f(z) = \sqrt{\frac{\hat{\sigma}_{zz}^2}{\hat{\sigma}_{zz}^2 + \bar{Z}^2}},$$

where  $\hat{\sigma}_{zz}^2 = \sum_{i=1}^n (Z_i - \bar{Z})^2/n$ ,  $\bar{Z} = \sum_{i=1}^n Z_i/n$ .

Note that we replace the unknown  $C$  by two components. First we use the consistent estimators  $\tilde{\sigma}_u$  and  $\hat{\sigma}_{zz}$ . Then for the correlation parameter we conduct a grid search, since we cannot estimate it. Therefore, if we know the true correlation (i.e.,  $\rho_0$ ), then:

$$\hat{C} = \tilde{\sigma}_u \hat{\sigma}_{zz} \rho_0 \xrightarrow{p} C = \sigma_u \sigma_{zz} \rho_0. \quad (5)$$

Of course if we do not know  $\rho_0$ , this will not be consistent. In the test statistics, theorems and the discussions below, (5) will be a good guide. An important point is to understand the relation between  $\rho_0$  and  $\beta_0$ . To that effect, see that by Assumption and (3)

$$\begin{aligned} EZ_i u_i &= C \\ EZ_i(y_i - X_i \beta_0) &= C \\ EZ_i y_i - (EZ_i X_i) \beta_0 &= \sigma_u \sigma_{zz} \rho_0. \end{aligned} \quad (6)$$

So clearly  $\beta_0, \rho_0$  are jointly unidentified. This also shows that  $\beta_0, \rho_0$  are linked. We also discuss how to build joint confidence intervals using (6) and test statistic here in subsection 3.5.

The key question is how do we obtain (4)? Why it is built in the way it is? This will be answered rigorously in the proof of Theorem 1, but here we provide a brief sketch. Lemma A.1i shows that bias in two-stage least squares estimate by using Assumption 1 is

$$(\pi_0^2 Q_{zz})^{-1} \pi_0 C,$$

when we have  $k = l = 1$ . If we know  $C$ , we can then subtract the least squares estimate of the bias from  $\hat{\beta} - \beta_0$  and setup the test. So set the test statistic at true  $C$  as

$$\frac{\sqrt{n}[\hat{\beta} - \beta_0 - (\hat{\pi}^2 \hat{Q}_{zz})^{-1} \hat{\pi} C]}{\tilde{\sigma}_u [\hat{\pi}^2 \hat{Q}_{zz}]^{-1/2}}.$$

Using (3) (5)

$$NT(\beta_0, \rho_0) = \frac{\sqrt{n}(\hat{\beta} - \beta_0)}{\hat{\sigma}_u |\hat{\pi}|^{-1} \hat{Q}_{zz}^{-1/2}} - \sqrt{n} \text{sgn}(\hat{\pi}) f(z) \rho_0. \quad (7)$$

Under the alternative of  $\rho = \rho_1 \neq \rho_0$ , NT statistic is evaluated at the wrong correlation between the instrument and the structural error,  $\rho_1$ . We denote the test statistic in that case as  $NT(\beta_0, \rho_1)$  to differentiate two cases of  $\rho_0$  (true correlation) and  $\rho_1$  (the wrong correlation choice). In the following Theorem, we consider the case of  $k = 1$  and  $l = k$  which is an empirically relevant case in most applied research. We analyze the null distribution in Theorem 1i, then in Theorem 1ii, we consider the limit of the modified test when we use the wrong correlation value.

**Theorem 1.** *Under Assumptions 1-2, with (3), when  $k = 1, l = k$ ,*

(i). *Under the null of  $H_0 : \beta = \beta_0, \rho = \rho_0$ ,*

$$NT(\beta_0, \rho_0) \xrightarrow{d} N(0, 1).$$

(ii). *Under the alternative  $\beta = \beta_0, \rho = \rho_1$ ,*

$$NT(\beta_0, \rho_1) \rightarrow \infty.$$

Remarks.

1. Theorem 1 shows that  $NT(\beta_0, \rho_0)$  converges to a standard normal limit if the true correlation is used (i.e.,  $\rho_0$ ) and if  $\beta = \beta_0$ . Otherwise the test diverges to infinity. Since the intention is to infer structural parameters we try a grid search over correlation values. This theorem can help applied researchers in their efforts for inference on structural parameters. Basically, in large samples if the null is true, then at the true correlation level we do not reject the null and all the other values of the correlation we reject the null hypotheses. We can have a very fine grid, and this helps us, as can be seen from Figures 1-2 in simulations.

An important issue in practice is the size distortion of the regular t-test due to the violation of exogeneity. Our test remedies this problem. The key issue is if the null is true, and  $H_0 : \beta = \beta_0, \rho \neq 0$ , can our test fail to reject the null? First of all, NT statistic at certain correlation values  $\rho \neq \rho_0$  can reject the true null. But since we are doing a grid search over correlation values at  $\rho = \rho_0 \neq 0$ , the test will not reject the null and have the correct inference on the structural parameters which sets  $\beta = \beta_0$ . The details of the grid over correlation values and empirics are shown further in this Remark, and Remark 2 below.

To be specific about the implementation of the test, we start with a reasonable value of the correlation between the instrument and the structural error. For example, we can start from -0.3. So at  $\rho = -0.3$  if the absolute value of  $NT(\beta_0, \rho_0)$  is larger than 1.96, then we do reject the null, and record this  $-0.3$  as not belonging to true correlation. We repeat this with a grid step of 0.01 in the positive direction. So say that at  $\rho = -0.1$ , the absolute value of  $NT(\beta_0, \rho_0)$  is less than 1.96, we record the correlation of -0.1 as belonging to true correlation set. Then we continue this process for other correlation values until maybe a reasonable upper bound is reached in correlation values. More about the details of this grid search is examined in Remark 2 below. The size and power issues will be discussed at length in the Remarks below.

2. What if we miss the true correlation in the grid search when the null hypotheses is true? We know that  $NT(\beta_0, \rho_1) \rightarrow \infty$ , and only if we know the true correlation  $\rho_0$  we do not reject the null if it is true. We can pinpoint where the true correlation lies by doing the grid search over the various correlation values and reporting the ones which fail to reject  $H_0$ .

Observe that the NT test is linear, continuous, and monotonic in the value of the correlation, which is clear from (4). In other words, when we start the grid search from -1, and go toward 1, the NT test will either decrease or increase depending on  $-sgn(\hat{\pi})$ . This is good news if we miss the true correlation in our grid search when the null is true.

To illustrate this point assume that the test is -1.96 at correlation 0, and 1.96 at correlation 0.1. Since the test is monotonic and continuous in correlation ( $\rho$ ), the NT statistic must be equal to each value between -1.96 and 1.96 for some correlation, because of the Intermediate Value Theorem. Therefore, in this example we know that the NT statistics that are evaluated at correlations between 0.0 and 0.1 are less than the 5% critical value, hence the null will not be rejected. So we can get the correct inference, and at the same time since the null is true, we can understand where the true correlation lies (i.e., in this example between 0 and 0.1). The power issue is analyzed in Remarks 6 and 7 below.

3. An important issue here is if the difference between the two-stage least squares estimate and  $\beta_0$  is positive and if the sign of the true correlation is the reverse of the sign of the first stage estimate, there is no need to be concerned about inference if we reject the null with regular t test, since the NT will be much larger (see that standard errors will be close since the bias is small). The same issue is true if the difference between the two-stage least squares estimate and  $\beta_0$  is negative and the sign of the correlation has the same sign of the first stage estimate then if we reject the null with regular t, we will reject with NT as well. So violation of exogeneity

will not change the results of the inference in this case as long as  $\hat{\sigma}_u$  is close to  $\tilde{\sigma}_u$ .

4. Another important point is the local analysis. Assume that  $\beta = \beta_0$  is true, and consider what will happen to the test statistic if we use  $n^{-1/2}$  neighborhood of  $\rho_0$ :  $\rho_0 + d/n^{1/2}$ , where  $d \neq 0$ . In other words, we make a minor mistake in the correlation choice. To state the case rigorously, we call this the local NT test ( $NT_l(\beta_0, \rho_0)$ )

$$\begin{aligned} NT_l(\beta_0, \rho_0) &= \frac{\sqrt{n}(\hat{\beta} - \beta_0)}{\hat{\sigma}_u[|\hat{\pi}|^{-1}\hat{Q}_{zz}^{-1/2}]} - \sqrt{n}\text{sgn}(\hat{\pi})f(z)(\rho_0 + d/n^{1/2}) \\ &= NT(\beta_0, \rho_0) - \text{sgn}(\hat{\pi})f(z)d \\ &\xrightarrow{d} N(D, 1), \end{aligned}$$

where we use (12), and  $D = -\text{sgn}(\hat{\pi})f(z)d$ .

So instead of  $N(0, 1)$  distribution as in  $NT(\beta_0, \rho_0)$  (where  $NT(\beta_0, \rho_0)$  assumes that we know the true correlation level) the true distribution is again normal with variance 1, but shifted to left or right. So this shows that there is local power when we choose the wrong correlation.

Then the question is: can we conduct inference in finite samples with such a minor violation? We answer this question in this remark and Remark 6 below. If  $\beta = \beta_0$ , then two things can happen. First, since we use wrong critical values (i.e.,  $N(0, 1)$  and the truth is  $N(D, 1)$ ), then  $NT_l(\beta_0, \rho_0)$  value may be not so large (compared with  $NT(\beta_0, \rho_0)$ ) and still we do not reject the null hypothesis of  $H_0 : \beta = \beta_0, \rho = \rho_0$ . So this is recorded as non rejection in our grid search of correlation values. In other words this may enlarge the true correlation set, and we fail to reject the null. But this is good in the sense that the truth is  $\beta = \beta_0$ , and our test still shows that result.

The second possibility is  $NT_l(\beta_0, \rho_0)$  is much larger than the  $N(0, 1)$  critical values and leads us to reject the null at that specific correlation level (i.e., at  $\rho_0 + d/n^{1/2}$ ). However, given Remark (2) of Theorem 1, we will detect the true correlation because of monotonicity and the intermediate value theorem, so this problem will not be relevant for inference on  $\beta = \beta_0$ .

5. If the alternative is true (i.e.,  $\beta_1$  is the true value and  $\beta_0 \neq \beta_1$ ), then clearly  $NT(\beta_1, \rho_0) \rightarrow \infty$  (when  $\rho = \rho_0$ ). There is power against the fixed alternatives for  $\beta$  at true value of the correlation (i.e.,  $\rho_0$ ).

As an additional fact, when  $H_0$  is false, and the true value of  $\beta$  is  $\beta_1$  and if we impose  $\beta_0$ , through Assumptions 1 and 2,

$$\tilde{\sigma}_u^2 - \sigma_u^2 \xrightarrow{p} a < \infty,$$

where  $a \neq 0$ ,

$$a = (\beta_1 - \beta_0)^2(\pi_0^2 Q_{zz} + \Sigma_{VV}) - 2(\beta_1 - \beta_0)(\pi_0 C + \Sigma_{uV}).$$

Under the alternative the  $\tilde{\sigma}_u^2$  is not consistent; however this does not affect the consistency of the NT statistic when we have fixed alternatives for  $\beta$ .

We now conduct another local power analysis. Set  $\beta_1 = \beta_0 + c/n^{1/2}$ ,  $c \neq 0$ , note that then  $a \rightarrow 0$ , so  $\tilde{\sigma}_u^2 \xrightarrow{p} \sigma_u^2$ .

The NT test at the true correlation

$$\begin{aligned} \frac{\sqrt{n}(\hat{\beta} - \beta_0)}{\tilde{\sigma}_u |\hat{\pi}|^{-1} \hat{Q}_{zz}^{-1}} &= \sqrt{n} \text{sgn}(\hat{\pi}) f(z) \rho_0 - \frac{\sqrt{n}(\beta_1 - \beta_0)}{\tilde{\sigma}_u |\hat{\pi}|^{-1} \hat{Q}_{zz}^{-1}} \\ &\xrightarrow{d} N(0, 1) - \frac{c}{\sigma_u |\pi_0|^{-1} Q_{zz}^{-1}} \\ &\equiv N(\tilde{c}, 1), \end{aligned}$$

where  $\tilde{c} = -c|\pi_0|Q_{zz}/\sigma_u$ .

So we have local power in NT test against local alternatives to  $\beta_0$ . This also shows through  $\tilde{c}$  that with strong instruments, the power will be large. Note that with strong instruments,  $\tilde{c}$  (a shift in the mean compared with standard normal) will be large and it will be easy to differentiate the alternative from the null.

6. Next we consider  $NT(\beta_1, \rho_1)$  and analyze whether it is plausible to fail to have a power loss. Below we show that this is probable at only implausibly large correlation values when we select strong instruments. The simulations also confirm this.

$$\begin{aligned} NT(\beta_1, \rho_1) &= \frac{\sqrt{n}(\hat{\beta} - \beta_1)}{\tilde{\sigma}_u [|\hat{\pi}|^{-1} \hat{Q}_{zz}^{-1/2}]} - \sqrt{n} \text{sgn}(\hat{\pi}) f(z) \rho_1 \\ &= \frac{\sqrt{n}(\hat{\beta} - \beta_0) - \sqrt{n}(\beta_1 - \beta_0)}{\tilde{\sigma}_u [|\hat{\pi}|^{-1} \hat{Q}_{zz}^{-1/2}]} - \sqrt{n} \text{sgn}(\hat{\pi}) f(z) \rho_0 - \sqrt{n} \text{sgn}(\hat{\pi}) f(z) (\rho_1 - \rho_0). \end{aligned}$$

It is possible then that

$$\frac{\sqrt{n}(\beta_1 - \beta_0)}{\tilde{\sigma}_u [|\hat{\pi}|^{-1} \hat{Q}_{zz}^{-1/2}]} \cong \sqrt{n} \text{sgn}(\hat{\pi}) f(z) (\rho_1 - \rho_0),$$

where these two terms may be equal to each other and cancel each other in the test statistic. Then  $NT(\beta_1, \rho_1)$  will not diverge to infinity but converge to a normal distribution. This may result in failure to reject the null. Now we show that with strong instruments, this issue may

occur at only implausibly large correlation values. By analyzing the left hand side of the above, and using Estimator of Concentration Parameter=  $CP : n\hat{\pi}^2\hat{Q}_{zz}/\hat{\sigma}_u^2$  we have that:

$$\sqrt{CP}(\beta_1 - \beta_0) \cong \sqrt{n}sgn(\hat{\pi})f(z)(\rho_1 - \rho_0). \quad (8)$$

So if the concentration parameter is large, then the possible non-rejection of the false null occurs at correlation values near -1 or +1. These are nearly implausible values in any given application (given that instruments are selected carefully, not randomly). So the problem can be avoided with large n or by using strong instruments.

7. Related to Remark 6 and Remark 2 above, we may have a non-rejection (of the null  $H_0$ ) region at certain correlation values if  $\rho_1 = \rho_0 + d/\sqrt{n}$ , and if the alternative hypotheses is true. This will not be a practical issue as we show. This is related to equation (8) above.

$$\sqrt{CP}(\beta_1 - \beta_0) \cong sgn(\hat{\pi})f(z)d. \quad (9)$$

But this may be avoided with large n or strong instruments, where the left hand and right hand sides will be far apart in (9).

From an applied perspective, if we combine the Remarks above, if the null is true, we have a range of correlation values that do not reject the null. Under the alternative, then we may avoid non-rejection with large samples or strong instruments or both. So in either case with our method we can conduct inference for structural parameters with correlation values different than 0. This is a significant improvement on the standard t test currently used.

### 3.2 The Comparison Between the Regular t and the NT

In the regular t test statistic, we consider  $H_0 : \beta = \beta_0$ . In NT we consider the joint null of  $H_0 : \beta = \beta_0, \rho = \rho_0$ . The joint null is needed, since inference on  $\beta$  depends on the correlation between the structural error and the instrument. The analysis applies to the overidentified case, but to make the comparison easier we demonstrate this using the just identified case. The standard t test, given ( $k = l = 1$ ), is:

$$t = \frac{\sqrt{n}(\hat{\beta} - \beta_0)}{\hat{\sigma}_u|\hat{\pi}|^{-1}\hat{Q}_{zz}^{-1/2}},$$

where  $\hat{\sigma}_u^2 = n^{-1} \sum_{i=1}^n (y_i - x_i\hat{\beta})^2$ .

The NT at true correlation is:

$$NT(\beta_0, \rho_0) = \frac{\sqrt{n}(\hat{\beta} - \beta_0)}{\tilde{\sigma}_u |\hat{\pi}|^{-1} \hat{Q}_{zz}^{-1/2}} - \sqrt{n} \text{sgn}(\hat{\pi}) f(z) \rho_0.$$

We refer to the test statistic evaluated at  $(\rho_1 \neq \rho_0)$  as  $NT(\beta_0, \rho_1)$ . The differences between the regular and NT are clear from the equations above. First, as discussed above,  $\hat{\sigma}_u \neq \tilde{\sigma}_u$ , and they are asymptotically equivalent only in the case of  $\hat{\beta} \xrightarrow{p} \beta_0$ . The second difference is the subtraction of the drift in the NT. The regular t specifically assumes that  $\rho_0 = 0$ , the NT does not assume that. When the null is true,  $\beta = \beta_0$  and  $\rho = \rho_0$ , our test will always fail to reject. On the other hand, if  $\beta = \beta_1 \neq \beta_0$ , finding the true correlation is not important, since NT is consistent at all correlation levels, both diverge to infinity and the test rejects.

Since the regular t test assumes  $\rho_0 = 0$ , if this is not true then under the null  $t \rightarrow \infty$ . This is illustrated in the simulation in Table 1. The size distortions with the regular t test are huge, and one can almost always reject the true null. The situation gets worse with larger sample sizes. In the  $NT(\beta_0, \rho_0)$  test,  $\rho$  does not have to be equal to zero, the test converges to standard normal distribution, and the test has excellent size (see Tables 2-4) at  $\rho_0$ .

Then the next question is, if  $\beta = \beta_0$ , what if there is a mistake in the true correlation choice in the NT test? In large samples, there are two possibilities, with a large mistake  $\rho_1 \neq \rho_0$ , the NT test ( $NT(\beta_0, \rho_1)$  in that case) diverges to infinity as shown in Theorem 1ii. If we have a fine grid search, we can catch the true correlation since NT values with the wrong correlation will be very high, and with the true correlation the test statistic will be between the critical values in the standard normal distribution. This point is also discussed in Remark 2 after Theorem 1. If the correlation is local to  $\rho_0$ , then the distribution is a normal distribution with drift, so we may reject the null or not depending on the magnitude of the drift. In the regular t ratio, if the true correlation is not 0, but local to some other number, then again the regular t diverges to infinity. So the regular t rejects  $\beta = \beta_0$  wrongly all the time. Only if the true correlation is 0 and we put  $\rho_1 = 0 + d/n^{1/2}$ , then the regular t has the same distribution as NT. This is the distribution in Remark 4 after Theorem 1, and Theorem 1 in Berkowitz, Caner, and Fang (2008).

Next, if the null is true, what can we say about the performance of regular t and the NT in finite samples? Here we compare them in a simulation. In Table 2, for  $n = 100$ , at 5% nominal size at  $\beta_0 = 0$ , regular t rejects the true null 20% at  $\rho_0 = 0.1$ , and 90% at  $\rho_0 = 0.3$ . In Table 4, at  $\beta_0 = 0$ , at true  $\rho_0 = 0.1$ ,  $NT(\beta_0, \rho_0)$  rejects at 5%, and at  $\rho_0 = 0.3$   $NT$  rejects at 3%. There

is still a very large difference between two. Even if we make a mistake in the choice of true correlation, still NT does better. For example, if we choose a correlation of 0 or 0.2, when the truth is 0.1 the NT rejects the true null at 16-17% compared with 20% rejection of the regular t. At true correlation of 0.3, if we make a mistake and use correlation of 0.2 or 0.4 in our test, the NT rejects at 14-15%, where as the regular t has 90% rejection rate of  $\beta_0 = 0$ .

Another issue is that if the regular t fails to reject the true null, is that true for the NT as well? In large samples, regular t test chooses the correct null only when  $\rho_0 = 0$ , this is true for  $NT(\beta_0, \rho_0)$  test as well, as is clear from Theorem 1i. In small samples, with  $n = 100$  and  $\rho_0 = 0$ , the size of  $NT(\beta_0, \rho_0)$  is 4.9% at the nominal 5% level (not shown in Tables). For standard t test, this is 5.3% as seen in Table 1.

When  $\beta = \beta_1 \neq \beta_0$ , both the standard t and the NT are consistent. The NT can have some power losses in finite samples but the discussion in Remarks 6-7 after Theorem 1 shows that this can be prevented through a choice of strong instruments. Clearly the performance of the NT statistic is far superior to the standard t test statistic.

### 3.3 The Overidentified Case of One Endogenous Regressor

In this case, since  $k = 1, l \geq k$ , we assume that

$$\text{corr}(Z_{im}, u_i) = \frac{\text{cov}(Z_{im}, u_i)}{\sigma_u \sqrt{\text{var}(Z_{im})}}, \quad (10)$$

for all  $m = 1, \dots, l$ . Using Assumption 1 we can rewrite (10) as, for all  $i = 1, \dots, n$ , at the true correlation

$$C_m = \sigma_u \sqrt{\text{var}(Z_{im})} \rho_0. \quad (11)$$

So with (11) we assume two things in addition to the first case analyzed in section 3.1. First, the instruments are such that  $\text{cov}(Z_{im}, Z_{ip}) = 0$  for all  $m \neq p, m = 1, \dots, l, p = 1, \dots, l, i = 1, 2, \dots, n$ . In other words, the instruments are not correlated with each other. In finite samples we can handle this through simple projections as discussed in Remark 2, after Theorem 2. The second assumption in (11) is for all instruments the true correlation between the structural error and the instrument is the same ( $\rho_0$ ). This is similar to regular instrumental variable estimation, where the claim is that the correlation between  $u_i$  and  $Z_{im}$  is the same (and 0) for all instruments. So here we extend this assumption to nonzero correlations. Our main results hold with different

correlations, but then multiple grid searches would be required. For the many instruments case, this may not be practical. We do not further consider that case here.

Now we construct the test statistic. Note that the former case, ( $k = l = 1$ ), is a special case of this more general formulation. The following test can be built using Lemma A.1i. The test statistic at the true C is:

$$\frac{\sqrt{n}(\hat{\beta} - \beta_0 - (\hat{\pi}'\hat{Q}_{zz}\hat{\pi})^{-1}\hat{\pi}'C)}{\tilde{\sigma}_u(\hat{\pi}'\hat{Q}_{zz}\hat{\pi})^{-1/2}}. \quad (12)$$

We can replace the infeasible test in (12) with the following by Assumption 1, (11), and extending (5) to a vector, at true value of the correlation ( $\rho_0$ ),

$$NT(\beta_0, \rho_0) = \frac{\sqrt{n}(\hat{\beta} - \beta_0 - (\hat{\pi}'\hat{Q}_{zz}\hat{\pi})^{-1}\hat{\pi}'[\tilde{\sigma}_u\sqrt{\widehat{var}(Z_1)}, \dots, \tilde{\sigma}_u\sqrt{\widehat{var}(Z_l)}]'\rho_0)}{\tilde{\sigma}_u(\hat{\pi}'\hat{Q}_{zz}\hat{\pi})^{-1/2}},$$

where  $\tilde{\sigma}_u$  is the square root of the estimator  $\tilde{\sigma}_u^2$ , and  $\widehat{var}(Z_m) = \frac{1}{n}\sum_{i=1}^n(Z_{im} - \bar{Z}_m)^2$  where  $\bar{Z}_m = n^{-1}\sum_{i=1}^n Z_{im}$  for  $m = 1, \dots, l$ . We can further simplify the test above as

$$NT(\beta_0, \rho_0) = \frac{\sqrt{n}(\hat{\beta} - \beta_0 - (\hat{\pi}'\hat{Q}_{zz}\hat{\pi})^{-1}\tilde{\pi}\rho_0)}{\tilde{\sigma}_u(\hat{\pi}'\hat{Q}_{zz}\hat{\pi})^{-1/2}}, \quad (13)$$

where  $\tilde{\pi} = \hat{\pi}'[\tilde{\sigma}_u\sqrt{\widehat{var}(Z_1)}, \dots, \tilde{\sigma}_u\sqrt{\widehat{var}(Z_l)}]'$  which is scalar.

Note that (12) simplifies to (4) as can be seen from (7). We test the joint null of  $H_0 : \beta = \beta_0, \rho = \rho_0$ .

**Theorem 2.** *Under Assumptions 1-2, with (11), and*

(i). *Under the null of  $H_0 : \beta = \beta_0, \rho = \rho_0$  when  $k = 1, l \geq k$*

$$NT(\beta_0, \rho_0) \xrightarrow{d} N(0, 1).$$

(ii). *Under alternative if  $\beta = \beta_0, \rho = \rho_1 \neq \rho_0$ , then*

$$NT(\beta_0, \rho_1) \rightarrow \infty.$$

Remarks.

1. Theorem 2 shows that  $NT(\beta_0, \rho_0)$  still has a standard normal distribution when  $k = 1, l \geq k$ . In large samples at the true correlation level the test statistic does not reject the null if  $H_0$  is true. At other values of correlation the test rejects the null. In the finite samples, this case is exactly the same as the just identified case. Choosing a fine grid with strong instruments ensures good size and power. If we evaluate at  $\rho_1 \neq \rho_0$  and still do not reject  $H_0$  (when  $\beta \neq \beta_0$

is false), then, as in the just identified case, choosing strong instruments solves the problem. If we evaluate at  $\rho_1 \neq \rho_0$ , then our test may reject the true null for that correlation, but this is easily fixed. Since the test is monotonic in the correlation, choosing a grid and conducting the tests in these new correlation values, we will be able to fail to reject the true null.

2. Note that in finite samples, instruments may be correlated. So we can use the following. Assume that we have two instruments:  $Z_{i1}, Z_{i2}$ , for  $i = 1, \dots, n$ . We regress (least squares)  $Z_{i1}$  on  $Z_{i2}$  and define the residual as  $Z_{i1\perp}$ . Then we use  $Z_{i1\perp}$  and  $Z_{i2}$  in the test.

### 3.4 The General Case

For testing individual coefficients when  $k > 1$  (multiple endogenous variables) and to get a consistent estimate for  $\sigma_u$  (i.e.,  $\tilde{\sigma}_u$ ) we need to impose  $\beta = \beta_0$  for all parameters. Then we have a joint NT test for  $H_0 : R\beta = R\beta_0, \rho = \rho_0$ , where  $R$  is a  $j \times k$  matrix. This test will include all parameters corresponding to endogenous regressors in the structural equation.

So we now introduce the NT evaluated at  $\beta_0, \rho_0$  (true correlation). This is constructed in the same way as before, and the proof is the same, and hence is skipped. Still we impose (11) with Assumption 1

$$C_m = \sigma_u \sqrt{\text{var} Z_{im} \rho_0}, \quad (14)$$

for all  $m = 1, \dots, l, i = 1, \dots, n$ .

We want to test  $H_0 : R\beta = R\beta_0, \rho = \rho_0$ . Define  $NT(\beta_0, \rho_0)$  as follows, by extending (5) to a vector

$$\begin{aligned} NT(\beta_0, \rho_0) &= n[R\hat{\beta} - R\beta_0 - R(\hat{\pi}'\hat{Q}_{zz}\hat{\pi})^{-1}\hat{\pi}'[\tilde{\sigma}_u\sqrt{\widehat{\text{var}}(Z_1)}, \dots, \tilde{\sigma}_u\sqrt{\widehat{\text{var}}(Z_l)}]'\rho_0]' \\ &\quad \times (R\tilde{\sigma}_u^2(\hat{\pi}'\hat{Q}_{zz}\hat{\pi})^{-1}R')^{-1} \\ &\quad \times [R\hat{\beta} - R\beta_0 - R(\hat{\pi}'\hat{Q}_{zz}\hat{\pi})^{-1}\hat{\pi}'[\tilde{\sigma}_u\sqrt{\widehat{\text{var}}(Z_1)}, \dots, \tilde{\sigma}_u\sqrt{\widehat{\text{var}}(Z_l)}]'\rho_0]. \end{aligned}$$

**Theorem 3.** *Under Assumptions 1-2, with (14), and*  
(i). *Under the joint null of  $R\beta = R\beta_0, \rho = \rho_0$ , we have*

$$NT(\beta_0, \rho_0) \xrightarrow{d} \chi_j^2.$$

(ii). *Under the alternative  $R\beta = R\beta_0, \rho = \rho_1 \neq \rho_0$ ,*

$$NT(\beta_0, \rho_1) \rightarrow \infty.$$

Remarks.

1. When we use  $\rho_1$  (make a mistake in selection of true correlation) then  $NT(\beta_0, \rho_1) \rightarrow \infty$ , as in Theorem 1 where  $\rho_1$  is used instead of  $\rho_0$ .

2. The test is consistent with a fixed alternative at  $\beta = \beta_1 \neq \beta_0$ . This can be shown easily. With a local alternative, also there is still power if we are at true correlation. When we use correlation values different than the true correlation  $\rho_1 \neq \rho_0$ , then it is possible to fail to reject the false null. However, as shown in Remarks 6-7 in the just identified case, using strong instruments solves the problem.

3. Note that with  $\beta = \beta_0$  and using a correlation value local to  $\rho_0$ , the test is distributed as noncentral  $\chi^2$  in large samples. So the issue is can we still make the correct inference about  $\beta = \beta_0$ ? As explained before, with a fine grid the NT will fail to reject at certain correlation values of the joint null of  $R\beta = R\beta_0, \rho = \rho_0$ .

### 3.5 Confidence Intervals

Theorems in the previous sections provide us with a way of building joint confidence interval for  $\beta_0, \rho_0$ . In the multivariate endogenous variables case, start by testing the null of  $H_0 : \beta = \beta_0, \rho = \rho_0$ . Then by special case of Theorem 3i (i.e.  $R = I_k$ , where  $I_k$  is the identity matrix of dimension  $k$ , so the limit in Theorem 3i is  $\chi_k^2$ ), we invert the test  $NT(\beta_0, \rho_0)$ , and get the joint confidence interval. So  $\{(\beta_0, \rho_0) : NT(\beta_0, \rho_0) \leq \chi_{k,\alpha}^2\}$  is an asymptotically valid 100  $(1 - \alpha)$  confidence set, where  $\chi_{k,\alpha}^2$  represents the 100  $\alpha$  critical value of the  $\chi_k^2$  distribution.

This also means that the test NT is inverted jointly on a grid search over  $\beta_0, \rho_0$  values. To warn the applied researchers we do not have a grid only on  $\rho_0$  and then use each value of  $\rho_0$  to find a confidence interval for  $\beta_0$  based on  $\hat{\beta}$  and standard errors. Rather, the grid search is joint over  $(\beta_0, \rho_0)$  values. Our test is asymptotically pivotal, and the joint confidence sets show that  $\beta_0$  can be unbounded with positive probability (if we have not normalized covariances as correlations in Assumption 1, we could have C 's unbounded in the joint confidence set as well). This can be seen from section 3 of Dufour (1997). So the confidence intervals are valid based on inverting NT statistic. We illustrate the construction of the joint confidence intervals in two empirical examples in Section 5.

### 3.6 Test for Exclusion Restriction

In this part of the paper, we propose a test for the exclusion restriction which can even be used in the just identified case. This is basically testing the null of  $H_0 : \beta = \beta_0, \rho = 0$ . So we benefit from  $NT(\beta_0, 0)$  where  $R = I_k$ . The test is

$$NT(\beta_0, 0) = \frac{n(\hat{\beta} - \beta_0)'(\hat{\pi}'\hat{Q}_{zz}\hat{\pi})(\hat{\beta} - \beta_0)}{\tilde{\sigma}_{uu}^2}.$$

The next Corollary derives the limit, and the following discussion shows how to implement the test to get a meaningful answer from the test for exclusion restrictions.

**Corollary 1.** *Under Assumptions 1-2, with (14), and under the joint null of  $\beta = \beta_0, \rho = 0$ , we have*

$$NT(\beta_0, 0) \xrightarrow{d} \chi_k^2.$$

Remarks.

1. The proof is a subcase of Theorem 3, and hence it is skipped.

2. The implementation is as follows. First setup a plausible grid values for  $\beta_0$ , a compact interval of  $[\beta_{0l}, \beta_{0u}]$ , where  $\beta_{0l}, \beta_{0u}$  represent the lower and upper bound of plausible  $\beta_0$  values. With these values we test  $NT(\beta_0, 0)$ . If for all plausible values the test rejects the null, then the instruments are not exogenous.

3. Note that this test is even valid for  $k = l$ , which is the just identified case. To our knowledge this is the first test statistic to do so in the literature.

4. The intuition of the test is such that by keeping  $\rho_0 = 0$ , and changing  $\beta_0$  values in the test, we assume if the model is true, we find such  $\beta_0$  and  $\rho_0 = 0$ , so the null will not be rejected. But if the instruments are not exogeneous, the test will be rejected at all grid values for  $\beta_0$ .

5. Regular t and Wald tests for testing only  $\beta = \beta_0$  are very similar in form, the difference being the  $\tilde{\sigma}_u^2$ . Previous literature such as Berkowitz, Caner and Fang (2008) already found that even with a minor violation of exogeneity, the distributions change and there is a drift which depends on the correlation between the instrument and the error. So in one sense, we are benefiting from that idea here. Instead of fixing  $\beta_0$  and to be tested once, we test a grid values of  $\beta_0$  in our test. If all of them reject the null then the instruments are not exogeneous.

## 4 Simulation

In this part of the paper we conduct simulations to answer the following questions. First, can we verify the results of Theorem 1? Namely, can we see that  $NT(\beta_0, \rho_0) \xrightarrow{d} N(0, 1)$ , and  $NT(\beta_0, \rho_1) \rightarrow \infty$  in large samples? The second issue is the finite sample behavior of the test statistic  $NT(\beta_0, \rho_0)$ . In the finite samples given a grid search (it may be a very fine grid search, with very small steps in a given empirical application), does the smallest rejection level still correspond to  $NT(\beta_0, \rho_0)$ ? The third question is related to power of the test. Is there a power loss at certain grid points as discussed after Theorem 1? If there is, can we also see that they are near extreme correlation values for a given application? If this power loss occurs away from  $[-0.3, 0.3]$  range of correlations, then that power loss may not be important as it will not arise in practice with reasonably chosen instruments. We generate the data with one instrument ( $l = 1$ ), one endogenous regressor ( $k = 1$ ) and no control variables.

$$y_i = X_i\beta_0 + u_i, \tag{15}$$

$$X_i = Z_i\pi_0 + V_i, \tag{16}$$

where  $\beta_0 = 0$ ,  $i = 1, \dots, n$ , and  $\pi_0 = 2$ . The structural error  $u_i$ , the first stage error  $V_i$ , and the instrument are iid. These are generated from the same joint normal distribution with  $N(0, \Lambda)$ , where

$$\Lambda = \begin{bmatrix} 1 & cov(Z_i, u_i) & 0 \\ cov(Z_i, u_i) & 1 & cov(V_i, u_i) \\ 0 & cov(u_i, V_i) & 1 \end{bmatrix},$$

since  $varZ_i = 1$ ,  $varu_i = 1$ ,  $cov(Z_i, u_i)$  is also the correlation between  $Z_i, u_i$ . This is denoted as  $\rho_0$  in the other sections. The covariance between  $V_i, u_i$  is set at 0.5. Since the variances are set at 1, the true correlation between the structural error and the instrument varies among -0.5, -0.3, -0.1, 0.1, 0.3, 0.5. The grid step is 0.1 for the tables. For the graphs the true correlation is set at -0.1 and the grid step is 0.01.<sup>1</sup> The sample sizes in the tables are  $n = 100, 200, 1000$ . The iteration number is 10000. For the size exercise, we report the percentage of rejections at 5% critical values from the standard normal distribution (-1.96, +1.96).

Table 2 provides the size of  $NT(\beta_0, \rho_0)$  and the rejection rate of the true null  $H_0 : \beta = 0, \rho = \rho_0$  for  $NT(\beta_0, \rho_1)$  (tests evaluated at  $\rho_1 \neq \rho_0$ ) at  $n = 1000$ . In Table 2,  $\rho_0$  represents the true

---

<sup>1</sup>The results look very similar when the true correlation is set at 0.25 or 0.52.

correlation between the structural error ( $u_i$ ) and the instrument ( $Z_i$ ). The first column is the grid values of the correlation “Grid”. When the grid value is equal to  $\rho_0$  then the size of the test should be 5% at that level ideally. Otherwise if the grid value of the correlation is not equivalent to  $\rho_0$  then the rejection rate of the test should be near 100% according to Theorem 1ii. We see that the results in Table 2 confirm Theorem 1. Namely, the size of the  $NT(\beta_0, \rho_0)$  test is at the 1-5% level (i.e., at  $\rho_0 = -0.5$ ,  $NT(\beta_0, \rho_0)$  is the one that corresponds to  $Grid = -0.5$ ). Otherwise when  $Grid = \rho_1 \neq \rho_0$  the test is  $NT(\beta_0, \rho_1)$ . When we look at the  $NT(\beta_0, \rho_1)$  test the rejection rate is 88-100% at the 5% nominal level. So if we have a grid search of the correlation, then only at the true value we will get the 5% rejection at nominal levels, otherwise we almost always reject the null. In that sense, we can differentiate the true correlation by looking at the absolute value of the NT statistic. If the absolute value of the test is less than 1.96 then the test fails to reject and the correlation that is used belongs to true correlation set.

To see how reliable this is in finite samples, we conduct the same exercise with  $n = 100, n = 200$  observations. In Tables 3 - 4 we see that  $NT(\beta_0, \rho_0)$  test achieves 1-5% size at 5% nominal level. This confirms that even in the finite samples the asymptotic approximation is very good. Table 3 shows the size of the test for  $NT(\beta_0, \rho_0)$  and the rejection rate of the true null for  $NT(\beta_0, \rho_1)$  at  $n = 200$ . For example, at true correlation of  $\rho_0 = -0.1$ , the  $NT(\beta_0, \rho_0)$  has the size of 4.5%, and  $NT(\beta_0, \rho_1)$  (which imposes  $\rho_1 = -0.2$ ) has the the rejection rate of 29.6% when the true null is  $\beta = 0, \rho = -0.1$ . But still there is substantial rejection rate difference between  $NT(\beta_0, \rho_1)$  and  $NT(\beta_0, \rho_0)$  tests. At  $n = 100$  in Table 4,  $NT(\beta_0, \rho_0)$  still has the smallest rejection rate. Tables 3-4 support our claim in Remarks 2 and 4 (in the just identified case) of the existence of a region of non-rejection of the true null when the null of  $\beta = 0, \rho = \rho_0$  is true. This region is around the true correlation value. We also report size results with a much finer grid of 0.01, these are shown in Figures 1 and 2, and verify Theorem 1.

Tables 5-8, report the percentage of the rejections when true  $\beta$  is not equal to zero. We have the same number of iterations as the size exercise, and the same critical values are used. The true values of  $\beta_0 = -2, -1, 1, 2$ , and we test  $H_0 : \beta = 0, \rho = \rho_0$ , and  $n = 100, 1000$ . The results confirm the remarks after Theorem 1. Namely, the power of  $NT$  is very good at almost all the relevant correlation levels for applications. Even at alternate correlation values (using  $\rho_1 = Grid \neq \rho_0$ ), the power is still very good in the range of the correlation values of  $[-0.3, 0.3]$ . There are certain power losses around highly implausible correlation levels, but, as can be seen, with large sample size this problem is less important. We also experiment with increasing the

concentration parameter estimate by putting  $\pi = 5$ , and this gives much better power results. We also experiment with  $\beta_0 = -0.5, -0.3, 0.3, 0.5$ , the results are very similar even in this close neighborhood of 0. These are not reported.

The applied researcher may use this method for a grid of correlations,  $\rho$ , between  $[-0.3, 0.3]$  for NT test. If one finds a region of non-rejection of the null, then this is also the confidence interval. If the grid search shows only rejection of the null, then the alternative is true. This test is a significant improvement to the regular t, which does not test the joint null and falsely rejects when  $\rho \neq 0$ .

In this simulation section, we also have size and power results for the exclusion restriction that is introduced. We test the joint null of  $H_0 : \beta = \beta_0, \rho = 0$ . We use the setup (15)(16), with the same covariance structure  $\Lambda$ . For the size part of the test we set  $\beta_0 = 0, \rho_0 = 0$ , and we evaluate  $NT(0, 0)$ . If the test is larger than the 5% critical value of  $\chi_1^2$  the we reject the null. There are 10000 iterations used. The sample size varies between 100, 200, 300, 1000. We record the percentage of the rejections of the true null. The results are in Table 9. It is clear that at all sample sizes the size is very good, and very close to 5%. So the exclusion restriction test has no size issue. The next issue is the power of the test. For this we propose two power tables. In this exercise (Table 10) the true model is  $\beta_0 = 0, \rho_0 \neq 0$ . However, the test imposes  $\beta = 0, \rho = 0$ . In this power exercise,  $\rho_0 = -0.5, -0.3, -0.1, 0.1, 0.3, 0.5$ . Again we record percentage of rejection of the null. This is done at 5% level for  $\chi_1^2$  distribution. We observe two things. If we have the correlation as -0.3 or 0.3, the test has very good power regardless of the sample size (87-100%). But if  $\rho_0 = 0.1$ , the exclusion restriction test has 16% power at  $n = 100$ . With  $n = 1000$ , the power increases to 88%. So large samples make a big difference if the true correlation is nonzero but close to zero. In the second power Table (Table 11), the true null is  $\rho_0 = 0$ , but  $\beta_0 \neq 0$ . And still we impose  $\beta = 0, \rho = 0$ . The results are similar to the Table 10. The power declines at  $n = 100$  when the structural parameter is nonzero but close to the zero, but with large samples, we can reject the null with probability one.

## 5 Implementation and Empirical Examples

The next step is to calculate  $NT(\beta_0, \rho_0)$  by iterating over potential values in a nested loop. To accomplish this, one calculates the test statistics over “reasonable” values of the correlation grid. Note that this could be the entire interval  $[-1, 1]$ , but, as discussed above, we restrict our

attention here to the range  $[-0.3, 0.3]$ . From the simulation results we find the statistic suffers from a loss of power at the extreme values of correlation. Since most instruments are carefully chosen,  $[-0.3, 0.3]$  is a reasonable range in practice. The range of  $\beta_0$  to test depends on the economic environment that dictates reasonable effect sizes and also what inferences the applied researcher chooses to draw.

If our interest centres on significance testing,  $H_0 : \beta_0 = 0, \rho = \rho_0$ , and if the applied researcher finds that the absolute value of  $NT$  drops below 1.96 in absolute value within a reasonable range of correlation values, then we fail to reject the null. On the other hand, if the researcher finds that the  $NT$  yields rejection at all reasonable levels of the grid, we can confidently reject the null and interpret the structural coefficient as being non-zero.

## 5.1 Empirical Examples

We apply our technique to two empirical examples. First, we replicate the results from Acemoglu and Johnson (2005), hereafter AJ. As discussed in the Introduction, the main results in AJ utilize early settler mortality to instrument for institutions when measuring the effect of institutions on economic growth as measured by GDP per capita. For our study, we have obtained the data used by AJ on 64 countries. In this discussion we focus on Table 2 of AJ which provides estimates for the just identified case of one instrument and one endogenous variable. In Table 2, Panel C, Column (3) of AJ, the two-stage least squares estimate of the effect of the constraint on executive power on GDP per capita is 0.756 with a standard error of 0.146. This coefficient is interpreted as highly statistically significant under standard inference.

Figure 3 presents a graphical depiction of the  $NT$  for reasonable grid values. The black region indicates combinations of  $\beta_0, \rho_0$  that cannot be rejected. This region can be interpreted as a confidence interval for the joint values of  $\beta_0, \rho_0$ . Notice that within the range  $[-0.30, +0.30]$   $NT$  indicates rejection for  $\beta_0 = 0$ . The regions of non-rejection are bounded away from  $\beta_0 = 0$  and demonstrate that, in fact,  $\beta_0 > 0$ . Note that even if  $\rho_0 \neq 0$  we can still infer that  $\beta > 0$ . Because the region of non-rejection includes  $\rho_0 = 0$ , the exclusion restriction test indicates that the we cannot reject perfect exogeneity in this example. Next we consider David Card's 1995 paper using proximity to a college as an instrument for educational attainment.<sup>2</sup> This paper

---

<sup>2</sup>The data set used in Card (1995) is available on David Card's website at: [http://emlab.berkeley.edu/users/card/data\\_sets.html](http://emlab.berkeley.edu/users/card/data_sets.html) (accessed May 19, 2009).

finds much larger returns to education relative to previous work. As in AJ, the instrumental variable in Card (1995) may not be completely exogenous, leading to biased results.

Figure 4 presents a graphical depiction of the  $NT$  zoomed in around the origin. Again, the black region indicates combinations of  $\beta_0, \rho_0$  that cannot be rejected. Note that in this case, the region of non-rejection is unbounded both above and below. Therefore, the confidence interval for  $\beta$  must include all possible values and no inference can be drawn. Notice again that the origin ( $\beta_0 = 0, \rho_0 = 0$ ) is not part of the non-rejected region. In fact,  $NT(0, 0) = 2.33$ . Under naive testing, this test statistic could be erroneously interpreted as indicating  $\beta \neq 0$ . But, as the figure clearly demonstrates, when considering the joint null, it is clear that without also considering  $\rho$ , one could draw incorrect inferences from rejection at  $\beta_0 = 0$ . We see clearly here that there is not enough information to identify  $\beta$  and we cannot reject that  $\rho$  is in the interval  $[-.1, .1]$ . The exclusion restriction test again indicates that for some values of  $\beta_0$  one cannot reject that  $\rho_0 = 0$ . Therefore, we cannot reject that the instrument is valid.

## 6 Discussion and Conclusion

In this paper we tackle one of the most important problems in applied econometrics, the violation of the exclusion restriction in instrumental variables (IV) estimation. We recognize that to infer anything about the structural parameters, one must know the correlation between the error and the instruments. Regular inference techniques assume that this correlation is zero. In any application this is a strong and likely incorrect assumption. We propose a joint test of structural parameters together with the correlation parameter. Since the correlation cannot be estimated in the test statistic that we develop, we conduct a grid search. This test corrects the bias in the two stage least squares estimators. Also we show that joint confidence intervals for  $\beta_0, \rho_0$  can be built by inverting this test statistic.

In future research, we want to extend these results to other contexts such as nonlinear GMM and possibly identification robust tests.

### REFERENCES

- Acemoglu, Daron, Simon Johnson, and James A. Robinson (2001), "The Colonial Origins of Comparative Development: An Empirical Investigation," *American Economic Review*, 91, 1369-1401.
- Acemoglu, Daron and Simon Johnson (2006), "Unbundling Institutions," *Journal of Political Economy*, 113, 949-995.

- Angrist, Joshua D. (1990), "Lifetime Earnings and the Vietnam Era Draft Lottery: Evidence from Social Security Administrative Records," *American Economic Review*, 80, 313-336.
- Angrist, Joshua D. and Alan B. Krueger (1991), "Does Compulsory School Attendance Affect Schooling and Earnings," *Quarterly Journal of Economics*, 106, 979-1014.
- Ashley, Richard, (2009), "Assessing the Credibility of Instrumental Variables Inference with Imperfect Instruments via Sensitivity Analysis," *Journal of Applied Econometrics*, Feb, 2009, Vol. 24: 325-337.
- Berkowitz, Daniel, Mehmet Caner, and Ying Fang (2008), "Are nearly exogenous instruments reliable?" *Economics Letters*, 101, 202-23.
- Berkowitz, Daniel, Mehmet Caner, and Ying Fang (2009), "The Validity of Instruments Revisited" Working paper, Dept. of Economics, North Carolina State University.
- Caner, Mehmet, (2009) "Near Exogeneity and Weak Identification in Generalized Empirical Likelihood Estimators: Many Moment Asymptotics," Working paper, Department of Economics, North Carolina State University.
- Card, David (1995), "Using Geographic Variation in College Proximity to Estimate the Returns to Schooling," in *Aspects of Labour Market Behaviour: Essays in Honor of John Vanderkamp*, eds. L. N. Christofides et al. Toronto: University of Toronto Press. 201-221.
- Djankov, Simeon, Rafael La Porta, Florencio Lopez-de-Silanes and Andrei Shleifer (2003), "Courts," *Quarterly Journal of Economics*, 118, 453-517.
- Duflo, Esther (2001), "Schooling and Labor Market Consequences of School Construction in Indonesia: Evidence from an Unusual Policy Experiment," *American Economic Review*, Vol. 91(4), pages 795-813, September.
- Dufour, Jean-Marie (1997), "Some Impossibility Theorems in Econometrics with Applications to Structural and Dynamic Models," *Econometrica*, vol.65, 1365-1389.
- Glaeser, Edward, Rafael La Porta, Florencio Lopez-de-Silanes and Andrei Shleifer (2004), "Do Institutions Cause Growth?" *Journal of Economic Growth*, 9, 271-303.
- Guggenberger, Patrik, (2009), "On the Asymptotic Size Distortion of tests when instruments locally violate the exogeneity assumption," Working paper, Department of Economics, UCLA.
- Guiso, Luigi, Paola Sapienza and Luigi Zingales (2006), "Does Culture Affect Economic Outcomes," *Journal of Economic Perspectives*, vol. 20(2), pages 23-48, Spring.
- Hall, Robert E. and Charles I. Jones (1999), "Why Do Some Countries Produce So Much More Output per Worker than Others?" *Quarterly Journal of Economics*, 114, 83-116.
- Kane, Thomas J. and Cecelia E. Rouse (1995), "Labor-Market Returns to Two- and Four-Year Colleges," *American Economic Review*, 85, 600-614.
- Kray, A. (2008) "Instrumental Variables Regression with Honestly Uncertain Exclusion Restrictions," unpublished manuscript. World Bank.
- Mauro, Paolo (1995), "Corruption and Growth," *Quarterly Journal of Economics*, 110, 681-712.
- Meghir, Costas and Marten Palme (2005), "Assessing the Effect of Schooling on Earnings Using a Social Experiment," *American Economic Review*, March 2005, pp 414-424.
- Nevo, Aviv, and Adam Rosen, (2008), "Identification with Imperfect Instrument," NBER Working Paper No. W14434, October 2008.

Figure 1: Simulation results for  $N=100$  when the correlation is  $-0.1$




Figure 2: Simulation results for  $N=1000$  when the correlation is  $-0.1$




Figure 3: Example Using Acemoglu and Johnson (2005)




Figure 4: Example Using Card (1995)




Table 1: Size (5% level), Standard t test

| $\rho_0 =$ | -0.5  | -0.3  | -0.1 | 0   | 0.1  | 0.3   | 0.5   |
|------------|-------|-------|------|-----|------|-------|-------|
| $n = 1000$ | 100.0 | 100.0 | 87.9 | 5.3 | 88.8 | 100.0 | 100.0 |
| $n = 200$  | 100.0 | 99.0  | 26.9 | 5.3 | 32.9 | 99.2  | 100.0 |
| $n = 100$  | 100.0 | 85.2  | 15.4 | 5.3 | 19.9 | 89.6  | 99.9  |

Note:  $\rho_0$  represents the true correlation between the single instrument and the error (second stage equation). The first column header is true correlation, all the other column headers are specific true correlation values.

Table 2: Rejection Rate of The Null  $\beta_0 = 0, \rho = \rho_0, NT(\beta_0, \rho_0), NT(\beta_0, \rho_1) n = 1000$

| Grid | $\rho_0 = -0.5$ | $\rho_0 = -0.3$ | $\rho_0 = -0.1$ | $\rho_0 = 0.1$ | $\rho_0 = 0.3$ | $\rho_0 = 0.5$ |
|------|-----------------|-----------------|-----------------|----------------|----------------|----------------|
| -1   | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.9 | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.8 | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.7 | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.6 | 94.7            | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.5 | 0.8             | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.4 | 94.1            | 90.8            | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.3 | 100.0           | 3.7             | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.2 | 100.0           | 90.0            | 88.7            | 100.0          | 100.0          | 100.0          |
| -0.1 | 100.0           | 100.0           | 4.8             | 100.0          | 100.0          | 100.0          |
| 0.0  | 100.0           | 100.0           | 88.7            | 88.1           | 100.0          | 100.0          |
| 0.1  | 100.0           | 100.0           | 100.0           | 4.9            | 100.0          | 100.0          |
| 0.2  | 100.0           | 100.0           | 100.0           | 88.2           | 90.4           | 100.0          |
| 0.3  | 100.0           | 100.0           | 100.0           | 100.0          | 3.1            | 100.0          |
| 0.4  | 100.0           | 100.0           | 100.0           | 100.0          | 91.4           | 94.4           |
| 0.5  | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 0.9            |
| 0.6  | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 95.1           |
| 0.7  | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| 0.8  | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| 0.9  | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| 1.0  | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |

Note: Grid represents the grid correlation values that we put into the NT tests. When  $Grid = \rho_0$ , the we have  $NT(\beta_0, \rho_0)$  test, otherwise the tests are  $NT(\beta_0, \rho_1)$ . The critical values are -1.96, +1.96. We set  $\pi_0 = 2$ . For example, in column 2,  $\rho_0 = -0.5$ , when  $Grid = -0.5$ , the test is called  $NT(\beta_0, \rho_0)$ , otherwise the tests are called  $NT(\beta_0, \rho_1)$ .

Table 3: Rejection Rate of The Null  $\beta_0 = 0, \rho = \rho_0, NT(\beta_0, \rho_0), NT(\beta_0, \rho_1)$   $n = 200$

| Grid | $\rho_0 = -0.5$ | $\rho_0 = -0.3$ | $\rho_0 = -0.1$ | $\rho_0 = 0.1$ | $\rho_0 = 0.3$ | $\rho_0 = 0.5$ |
|------|-----------------|-----------------|-----------------|----------------|----------------|----------------|
| -1   | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.9 | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.8 | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.7 | 88.8            | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.6 | 24.0            | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.5 | 1.1             | 83.4            | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.4 | 22.7            | 27.0            | 99.0            | 100.0          | 100.0          | 100.0          |
| -0.3 | 86.5            | 3.1             | 80.9            | 100.0          | 100.0          | 100.0          |
| -0.2 | 100.0           | 26.4            | 28.9            | 98.9           | 100.0          | 100.0          |
| -0.1 | 100.0           | 82.8            | 4.9             | 80.7           | 100.0          | 100.0          |
| 0.0  | 100.0           | 100.0           | 29.5            | 28.0           | 100.0          | 100.0          |
| 0.1  | 100.0           | 100.0           | 80.8            | 4.5            | 82.1           | 100.0          |
| 0.2  | 100.0           | 100.0           | 98.8            | 29.1           | 27.5           | 100.0          |
| 0.3  | 100.0           | 100.0           | 100.0           | 81.4           | 3.0            | 86.2           |
| 0.4  | 100.0           | 100.0           | 100.0           | 100.0          | 27.8           | 22.4           |
| 0.5  | 100.0           | 100.0           | 100.0           | 100.0          | 83.4           | 1.0            |
| 0.6  | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 24.3           |
| 0.7  | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 89.2           |
| 0.8  | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| 0.9  | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| 1.0  | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |

Note: Grid represents the grid correlation values that we put into the NT tests. When  $Grid = \rho_0$ , the we have  $NT(\beta_0, \rho_0)$  test, otherwise the tests are  $NT(\beta_0, \rho_1)$ . The critical values are -1.96, +1.96. We set  $\pi = 2$ . For example, in column 2,  $\rho_0 = -0.5$ , when  $Grid = -0.5$ , the test is called  $NT(\beta_0, \rho_0)$ , otherwise the tests are called  $NT(\beta_0, \rho_1)$ .

Table 4: Rejection Rate of The Null  $\beta_0 = 0, \rho = \rho_0, NT(\beta_0, \rho_0), NT(\beta_0, \rho_1)$   $n = 100$

| Grid | $\rho_0 = -0.5$ | $\rho_0 = -0.3$ | $\rho_0 = -0.1$ | $\rho_0 = 0.1$ | $\rho_0 = 0.3$ | $\rho_0 = 0.5$ |
|------|-----------------|-----------------|-----------------|----------------|----------------|----------------|
| -1   | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.9 | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.8 | 93.5            | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.7 | 52.8            | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.6 | 11.2            | 88.5            | 100.0           | 100.0          | 100.0          | 100.0          |
| -0.5 | 1.0             | 51.5            | 98.3            | 100.0          | 100.0          | 100.0          |
| -0.4 | 8.1             | 15.8            | 85.7            | 100.0          | 100.0          | 100.0          |
| -0.3 | 51.7            | 2.7             | 51.9            | 97.9           | 100.0          | 100.0          |
| -0.2 | 90.3            | 13.6            | 16.6            | 85.1           | 100.0          | 100.0          |
| -0.1 | 100.0           | 52.1            | 4.5             | 51.6           | 98.5           | 100.0          |
| 0.0  | 100.0           | 87.0            | 16.5            | 16.2           | 86.6           | 100.0          |
| 0.1  | 100.0           | 98.5            | 51.3            | 4.6            | 52.4           | 100.0          |
| 0.2  | 100.0           | 100.0           | 85.4            | 16.8           | 13.8           | 90.1           |
| 0.3  | 100.0           | 100.0           | 97.9            | 51.2           | 2.9            | 51.7           |
| 0.4  | 100.0           | 100.0           | 100.0           | 85.7           | 14.9           | 8.1            |
| 0.5  | 100.0           | 100.0           | 100.0           | 98.4           | 52.2           | 1.2            |
| 0.6  | 100.0           | 100.0           | 100.0           | 100.0          | 88.4           | 11.7           |
| 0.7  | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 52.5           |
| 0.8  | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 93.7           |
| 0.9  | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |
| 1.0  | 100.0           | 100.0           | 100.0           | 100.0          | 100.0          | 100.0          |

Note: Grid represents the grid correlation values that we put into the NT tests. When  $Grid = \rho_0$ , the we have  $NT(\beta_0, \rho_0)$  test, otherwise the tests are  $NT(\beta_0, \rho_1)$ . The critical values are -1.96, +1.96. We set  $\pi = 2$ . For example, in column 2,  $\rho_0 = -0.5$ , when  $Grid = -0.5$ , the test is called  $NT(\beta_0, \rho_0)$ , otherwise the tests are called  $NT(\beta_0, \rho_1)$ .

Table 5: Rejection percentage of  $H_0 : \beta = 0, \rho_0 = 0.1, NT(\beta, \rho_0), NT(\beta, \rho_1)$

| Grid | $n = 1000$     |                |               |               | $n = 100$      |                |               |               |
|------|----------------|----------------|---------------|---------------|----------------|----------------|---------------|---------------|
|      | $\beta_0 = -2$ | $\beta_0 = -1$ | $\beta_0 = 1$ | $\beta_0 = 2$ | $\beta_0 = -2$ | $\beta_0 = -1$ | $\beta_0 = 1$ | $\beta_0 = 2$ |
| -1   | 0.0            | 100.0          | 100.0         | 100.0         | 0.0            | 0.0            | 100.0         | 100.0         |
| -0.9 | 0.0            | 0.0            | 100.0         | 100.0         | 0.0            | 0.0            | 100.0         | 100.0         |
| -0.8 | 0.0            | 100.0          | 100.0         | 100.0         | 0.0            | 0.0            | 100.0         | 100.0         |
| -0.7 | 79.2           | 100.0          | 100.0         | 100.0         | 79.2           | 24.9           | 100.0         | 100.0         |
| -0.6 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.5 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.4 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.3 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.2 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.1 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.0  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.1  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.2  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.3  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.4  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.5  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 94.7          | 100.0         |
| 0.6  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 25.1          | 90.1          |
| 0.7  | 100.0          | 100.0          | 76.6          | 100.0         | 100.0          | 100.0          | 0.0           | 0.7           |
| 0.8  | 100.0          | 100.0          | 0.8           | 0.5           | 100.0          | 100.0          | 0.1           | 0.0           |
| 0.9  | 100.0          | 100.0          | 100.0         | 41.1          | 100.0          | 100.0          | 7.2           | 0.1           |
| 1.0  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 81.3          | 14.4          |

Note: Grid represents the grid correlation values that we put into the NT tests. When  $\rho_0 = Grid$ , then we have  $NT(\beta, \rho_0)$  test, otherwise the tests are  $NT(\beta, \rho_1)$ . The critical values are -1.96, +1.96. We set  $\pi_0 = 2$ . Here  $NT(\beta, \rho_0)$  is when  $\rho_0 = Grid = 0.1$ .

Table 6: Rejection percentage of  $H_0 : \beta = 0, \rho_0 = -0.1, NT(\beta, \rho_0), NT(\beta, \rho_1)$

| Grid | $n = 1000$     |                |               |               | $n = 100$      |                |               |               |
|------|----------------|----------------|---------------|---------------|----------------|----------------|---------------|---------------|
|      | $\beta_0 = -2$ | $\beta_0 = -1$ | $\beta_0 = 1$ | $\beta_0 = 2$ | $\beta_0 = -2$ | $\beta_0 = -1$ | $\beta_0 = 1$ | $\beta_0 = 2$ |
| -1   | 0.0            | 100.0          | 100.0         | 100.0         | 0.0            | 100.0          | 100.0         | 100.0         |
| -0.9 | 0.0            | 0.0            | 100.0         | 100.0         | 0.0            | 74.3           | 100.0         | 100.0         |
| -0.8 | 0.0            | 100.0          | 100.0         | 100.0         | 0.0            | 13.3           | 100.0         | 100.0         |
| -0.7 | 89.5           | 100.0          | 100.0         | 100.0         | 90.3           | 0.5            | 100.0         | 100.0         |
| -0.6 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 0.3            | 100.0         | 100.0         |
| -0.5 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 31.2           | 100.0         | 100.0         |
| -0.4 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 89.5           | 100.0         | 100.0         |
| -0.3 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.2 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.1 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.0  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.1  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.2  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.3  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.4  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.5  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 80.7          | 100.0         |
| 0.6  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 7.7           | 80.2          |
| 0.7  | 100.0          | 100.0          | 5.4           | 100.0         | 100.0          | 100.0          | 0.0           | 0.2           |
| 0.8  | 100.0          | 100.0          | 45.9          | 0.0           | 100.0          | 100.0          | 0.6           | 0.0           |
| 0.9  | 100.0          | 100.0          | 100.0         | 86.0          | 100.0          | 100.0          | 24.3          | 0.1           |
| 1.0  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 95.3          | 26.8          |

Note: Grid represents the grid correlation values that we put into the NT tests. When  $\rho_0 = Grid$ , then we have  $NT(\beta, \rho_0)$  test, otherwise the tests are  $NT(\beta, \rho_1)$ . The critical values are -1.96, +1.96. We set  $\pi_0 = 2$ . Here  $NT(\beta, \rho_0)$  is when  $\rho_0 = Grid = -0.1$ .

Table 7: Rejection percentage of  $H_0 : \beta = 0, \rho_0 = 0.3, NT(\beta, \rho_0), NT(\beta, \rho_1)$

| Grid | $n = 1000$     |                |               |               | $n = 100$      |                |               |               |
|------|----------------|----------------|---------------|---------------|----------------|----------------|---------------|---------------|
|      | $\beta_0 = -2$ | $\beta_0 = -1$ | $\beta_0 = 1$ | $\beta_0 = 2$ | $\beta_0 = -2$ | $\beta_0 = -1$ | $\beta_0 = 1$ | $\beta_0 = 2$ |
| -1   | 100.0          | 100.0          | 100.0         | 100.0         | 0.0            | 1.3            | 100.0         | 100.0         |
| -0.9 | 0.0            | 0.0            | 100.0         | 100.0         | 0.0            | 0.0            | 100.0         | 100.0         |
| -0.8 | 100.0          | 90.1           | 100.0         | 100.0         | 0.0            | 0.0            | 100.0         | 100.0         |
| -0.7 | 100.0          | 100.0          | 100.0         | 100.0         | 6.7            | 9.9            | 100.0         | 100.0         |
| -0.6 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.5 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.4 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.3 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.2 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.1 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.0  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.1  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.2  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.3  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.4  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.5  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.6  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 55.1          | 95.7          |
| 0.7  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 0.0           | 2.5           |
| 0.8  | 100.0          | 100.0          | 0.0           | 3.1           | 100.0          | 100.0          | 0.0           | 0.0           |
| 0.9  | 100.0          | 100.0          | 100.0         | 41.8          | 100.0          | 100.0          | 1.0           | 0.0           |
| 1.0  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 53.4          | 6.8           |

Note: Grid represents the grid correlation values that we put into the NT tests. When  $\rho_0 = Grid$ , then we have  $NT(\beta, \rho_0)$  test, otherwise the tests are  $NT(\beta, \rho_1)$ . The critical values are -1.96, +1.96. We set  $\pi_0 = 2$ . Here  $NT(\beta, \rho_0)$  is when  $\rho_0 = Grid = 0.3$ .

Table 8: Rejection percentage of  $H_0 : \beta = 0, \rho_0 = -0.3$ ,  $NT(\beta, \rho_0)$ ,  $NT(\beta, \rho_1)$

| Grid | $n = 1000$     |                |               |               | $n = 100$      |                |               |               |
|------|----------------|----------------|---------------|---------------|----------------|----------------|---------------|---------------|
|      | $\beta_0 = -2$ | $\beta_0 = -1$ | $\beta_0 = 1$ | $\beta_0 = 2$ | $\beta_0 = -2$ | $\beta_0 = -1$ | $\beta_0 = 1$ | $\beta_0 = 2$ |
| -1   | 98.6           | 100.0          | 100.0         | 100.0         | 0.0            | 0.0            | 100.0         | 100.0         |
| -0.9 | 0.0            | 0.0            | 100.0         | 100.0         | 0.0            | 0.0            | 100.0         | 100.0         |
| -0.8 | 100.0          | 100.0          | 100.0         | 100.0         | 0.0            | 0.0            | 100.0         | 100.0         |
| -0.7 | 100.0          | 100.0          | 100.0         | 100.0         | 96.5           | 91.8           | 100.0         | 100.0         |
| -0.6 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.5 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.4 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.3 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.2 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| -0.1 | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.0  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.1  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.2  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.3  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 100.0         |
| 0.4  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 97.2          | 100.0         |
| 0.5  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 56.4          | 100.0         |
| 0.6  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 1.5           | 67.3          |
| 0.7  | 100.0          | 100.0          | 0.0           | 100.0         | 100.0          | 100.0          | 0.1           | 0.0           |
| 0.8  | 100.0          | 100.0          | 98.3          | 0.0           | 100.0          | 100.0          | 3.7           | 0.0           |
| 0.9  | 100.0          | 100.0          | 100.0         | 98.9          | 100.0          | 100.0          | 50.1          | 0.4           |
| 1.0  | 100.0          | 100.0          | 100.0         | 100.0         | 100.0          | 100.0          | 100.0         | 40.7          |

Note: Grid represents the grid correlation values that we put into the NT tests. When  $Grid = \rho_0$ , then we have  $NT(\beta, \rho_0)$  test, otherwise the tests are  $NT(\beta, \rho_1)$ . The critical values are -1.96, +1.96.  $\pi_0 = 2$ . Here  $NT(\beta, \rho_0)$  is when  $Grid = \rho_0 = -0.3$ .

Table 9: Size (5% level), The Exclusion Restrictions Test ( $\beta_0 = 0, \rho_0 = 0$ )

| $n = 100$ | $n = 200$ | $n = 300$ | $n = 1000$ |
|-----------|-----------|-----------|------------|
| 4.6       | 5.0       | 4.9       | 5.0        |

Note: We impose  $\beta_0 = 0, \rho_0 = 0$  in the test, and compare with 5% critical values of  $\chi_1^2$ . The number of iterations are 10000.

Table 10: Power of the Exclusion Restriction Test (True Model:  $\beta_0 = 0, \rho = \rho_0$ )

| $\rho_0 =$ | -0.5  | -0.3  | -0.1 | 0.1  | 0.3   | 0.5   |
|------------|-------|-------|------|------|-------|-------|
| $n = 100$  | 100.0 | 87.0  | 16.5 | 16.2 | 86.6  | 100.0 |
| $n = 200$  | 100.0 | 100.0 | 29.5 | 28.0 | 100.0 | 100.0 |
| $n = 1000$ | 100.0 | 100.0 | 88.7 | 88.1 | 100.0 | 100.0 |

Note: We impose  $\beta_0 = 0, \rho = 0$ , and the true model is  $\beta_0 = 0, \rho_0 \neq 0$ . The details are given in the note to Table 9.

Table 11: Power of the Exclusion Restriction Test (True Model:  $\beta = \beta_0, \rho = 0$ )

| $\beta_0 =$ | -2    | -1    | -0.2  | 0.2   | 1     | 2     |
|-------------|-------|-------|-------|-------|-------|-------|
| $n = 100$   | 100.0 | 100.0 | 57.8  | 42.8  | 99.9  | 100.0 |
| $n = 200$   | 100.0 | 100.0 | 86.5  | 71.5  | 100.0 | 100.0 |
| $n = 1000$  | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 |

Note: We impose  $\beta = 0, \rho = 0$ , and the true model is  $\beta_0 \neq 0, \rho_0 = 0$ . The details are given in the note to Table 9.

## Appendix A: Proofs

Before the proof of the main Theorem, we need the following result that will help us in understanding the main result. This holds for both the simple case just identified case,  $k = l = 1$ , and the overidentified case  $k = 1, l > k$ .

**Lemma A.1.** *Under Assumptions 1-2, and under the null hypotheses of  $H_0 : \beta = \beta_0, \rho = \rho_0$ ,*  
(i).

$$\hat{\beta} - \beta_0 \xrightarrow{p} [\pi_0' Q_{zz} \pi_0]^{-1} [\pi_0' C] \neq 0.$$

(ii).

$$\hat{\sigma}_u^2 \xrightarrow{p} \sigma_u^2 - 2C_\pi(\pi_0' C + \Sigma_{Vu}) + C_\pi^2(\pi_0' Q_{zz} \pi_0 + \Sigma_{VV}).$$

(iii).

$$t_{2sls} \rightarrow \infty,$$

where  $t_{2sls}$  represents the regular two stage least squares based  $t$ -test.

**Proof of Lemma A.1.** We analyze a system with  $k = 1$ , and multiple instruments ( $l \geq k$ ). First we show that  $\hat{\beta}$  is inconsistent given Assumption 1.

$$\hat{\beta} - \beta_0 = \left[ \left( \frac{X'Z}{n} \right) \left( \frac{Z'Z}{n} \right)^{-1} \left( \frac{Z'X}{n} \right) \right]^{-1} \left[ \left( \frac{X'Z}{n} \right) \left( \frac{Z'Z}{n} \right)^{-1} \left( \frac{Z'u}{n} \right) \right]. \quad (17)$$

See that by first stage equation and Assumption 2

$$\begin{aligned} \frac{Z'X}{n} &= \frac{Z'Z}{n} \pi_0 + \frac{Z'V}{n} \\ &\xrightarrow{p} Q_{zz} \pi_0. \end{aligned} \quad (18)$$

$$n^{-1} \sum_{i=1}^n Z_i u_i \xrightarrow{p} E Z_i u_i = C < \infty, \quad (19)$$

where note the transformation in (5). Use (18)(19) in (17) to have

$$\hat{\beta} - \beta_0 \xrightarrow{p} [\pi_0' Q_{zz} \pi_0]^{-1} [\pi_0' C] \neq 0, \quad (20)$$

as long as  $C \neq 0$ . Next we show that  $\hat{\sigma}_u^2$  is not a consistent estimator for  $\sigma_u^2$ . First

$$\begin{aligned} n^{-1} \sum_{i=1}^n (y_i - x_i' \hat{\beta})^2 &= n^{-1} \sum_{i=1}^n (u_i - (x_i' \hat{\beta} - \beta_0))^2 \\ &= n^{-1} \sum_{i=1}^n u_i^2 - 2(\hat{\beta} - \beta_0) n^{-1} \sum_{i=1}^n x_i u_i \\ &\quad + (\hat{\beta} - \beta_0)' n^{-1} \sum_{i=1}^n x_i x_i'. \end{aligned} \quad (21)$$

See that from (20), set  $C_\pi = [\pi'_0 Q_{zz} \pi_0]^{-1} [\pi'_0 C]$ , by Assumption 2i, Assumption 1, and using the first stage equation

$$\begin{aligned} \frac{X'u}{n} &= \frac{\pi'_0 Z'u}{n} + \frac{V'u}{n} \\ &\xrightarrow{p} \pi'_0 C + \Sigma_{Vu}, \end{aligned} \quad (22)$$

$$\begin{aligned} \frac{X'X}{n} &= \frac{\pi'_0 Z'Z \pi_0}{n} + \frac{2\pi'_0 Z'V}{n} + \frac{V'V}{n} \\ &\xrightarrow{p} \pi'_0 Q_{zz} \pi_0 + \Sigma_{VV}. \end{aligned} \quad (23)$$

Use (20)(22)(23) in (21)

$$\hat{\sigma}_u^2 \xrightarrow{p} \sigma_u^2 - 2C_\pi(\pi'_0 C + \Sigma_{Vu}) + C_\pi^2(\pi'_0 Q_{zz} \pi_0 + \Sigma_{VV}) < \infty. \quad (24)$$

So the last two terms are nonzero (unless they cancel each other in special empirical cases). We cannot use  $\hat{\sigma}_u^2$  as a consistent estimator. Next we show that under Assumption 1, the t-test for  $H_0 : \beta = \beta_0, \rho = \rho_0$

$$t_{2sls} = \frac{\sqrt{n}(\hat{\beta} - \beta_0)}{\hat{\sigma}_u (X'P_Z X)^{-1/2}} \rightarrow \infty,$$

by (17)(20)(24). **Q.E.D.**

**Proof of Theorem 1.** The proof is a subcase of the proof of Theorem 2, since that proof is for  $k = 1, l \geq k$ .

**Proof of Theorem 2.** This proof is for  $k = 1, l \geq k$ , and hence covers the cases of  $k = l = 1$ , and  $k = 1, l > k$ . Now we show that NT test converges in distribution to standard normal distribution. In that respect, we first try to understand the numerator of the new test statistic. See that

$$\begin{aligned} n^{1/2}(\hat{\beta} - \beta_0) &= \left[ \left( \frac{X'Z}{n} \right) \left( \frac{Z'Z}{n} \right)^{-1} \left( \frac{Z'X}{n} \right) \right]^{-1} \\ &\times \left( \frac{X'Z}{n} \right) \left( \frac{Z'Z}{n} \right)^{-1} \left( \frac{1}{n^{1/2}} \sum_{i=1}^n (Z_i u_i - E Z_i u_i) + n^{1/2} E Z_i u_i \right) \\ &= \left\{ \left[ \left( \frac{X'Z}{n} \right) \left( \frac{Z'Z}{n} \right)^{-1} \left( \frac{Z'X}{n} \right) \right]^{-1} \left( \frac{X'Z}{n} \right) \left( \frac{Z'Z}{n} \right)^{-1} \left( \frac{1}{n^{1/2}} \sum_{i=1}^n (Z_i u_i - E Z_i u_i) \right) \right\} \\ &+ \left\{ \left[ \left( \frac{X'Z}{n} \right) \left( \frac{Z'Z}{n} \right)^{-1} \left( \frac{Z'X}{n} \right) \right]^{-1} \left( \frac{X'Z}{n} \right) \left( \frac{Z'Z}{n} \right)^{-1} \sqrt{n} C \right\} \\ &= A_1 + A_2, \end{aligned} \quad (25)$$

where  $A_1, A_2$  represent the first and the second terms with curly bracket expressions. By using Assumption 2, (18)

$$A_1 \xrightarrow{d} N(0, \sigma_u^2(\pi_0' Q_{zz} \pi_0)^{-1}). \quad (26)$$

Then by (18) and Assumptions 1, 2ii

$$A_2 \rightarrow \infty.$$

So we definitely have to subtract  $A_2$  from  $\sqrt{n}(\hat{\beta} - \beta_0)$  term. But the real issue is the handling of C. So we handle that by the arguments in the main text. Given (18)(25)(26) we have

$$NT(\beta_0, \rho_0) = \frac{\sqrt{n}(\hat{\beta} - \beta_0 - (\hat{\pi}' \hat{Q}_{zz} \hat{\pi})^{-1} \hat{\pi}' C)}{\tilde{\sigma}_u(\hat{\pi}' \hat{Q}_{zz} \hat{\pi})^{-1/2}} \xrightarrow{d} N(0, 1). \quad (27)$$

Equivalently via (12)(13), by writing the NT test in (27) as

$$NT(\beta_0, \rho_0) = \frac{\sqrt{n}(\hat{\beta} - \beta_0 - (\hat{\pi}' \hat{Q}_{zz} \hat{\pi})^{-1} \tilde{\pi} \rho_0)}{\tilde{\sigma}_u(\hat{\pi}' \hat{Q}_{zz} \hat{\pi})^{-1/2}}.$$

So under the joint null of  $\beta = \beta_0, \rho = \rho_0$ , then

$$NT(\beta_0, \rho_0) \xrightarrow{d} N(0, 1),$$

as shown above.

If we had used  $\rho_1 \neq \rho_0$  in our grid search

$$\begin{aligned} NT(\beta_0, \rho_1) &= \frac{\sqrt{n}(\hat{\beta} - \beta_0 - (\hat{\pi}' \hat{Q}_{zz} \hat{\pi})^{-1} \tilde{\pi} \rho_1)}{\tilde{\sigma}_u(\hat{\pi}' \hat{Q}_{zz} \hat{\pi})^{-1/2}} \\ &= \frac{\sqrt{n}(\hat{\beta} - \beta_0 - (\hat{\pi}' \hat{Q}_{zz} \hat{\pi})^{-1} \tilde{\pi} \rho_0)}{\tilde{\sigma}_u(\hat{\pi}' \hat{Q}_{zz} \hat{\pi})^{-1/2}} \\ &\quad + \frac{\sqrt{n}(\hat{\pi}' \hat{Q}_{zz} \hat{\pi})^{-1} \tilde{\pi}(\rho_0 - \rho_1)}{\tilde{\sigma}_u(\hat{\pi}' \hat{Q}_{zz} \hat{\pi})^{-1/2}} \\ &= NT(\beta_0, \rho_0) + \frac{(\hat{\pi}' \hat{Q}_{zz} \hat{\pi})^{-1} \tilde{\pi} n^{1/2}(\rho_0 - \rho_1)}{\tilde{\sigma}_u(\hat{\pi}' \hat{Q}_{zz} \hat{\pi})^{-1/2}} \\ &\rightarrow \infty. \end{aligned} \quad (28)$$

Note that the second term on the right hand side of the above equation diverges to infinity since  $\sqrt{n}(\rho_1 - \rho_0) \rightarrow \infty$ . **Q.E.D.**

## Appendix B: Stata Code

In this code,  $x$  is the endogenous variable of interest,  $y$  is the dependent variable, and  $z$  is the instrument. For simplicity, let 'covars' be a local macro for all exogenous covariates.

```
% ivreg y (x = z) 'covars'
% scalar b2sls = _b[x]
% scalar N = e(N)

% reg x 'covars'
% predict xresid, resid

% reg y 'covars'
% predict yresid, resid

% reg z 'covars'
% predict zresid, resid

% reg xresid yresid
% scalar phihat = _b[zresid]

% egen qzz = sum(zresid^2/N)

Choose a value of the structural parameter to test or do a grid search:
% forvalues b0 = -10(.01)10{

% egen ssyresid = total((yresid - xresid*b0)^2)
% scalar sigmatilda2 = 1/(N-1)*ssyresid;
% scalar sigmatilda = sqrt(sigmatilda2);

Choose a correlation equal to some value 'corr' or do a grid search:
% forvalues rho0 = -1(.01)1{
% scalar neat = N^(1/2)*(b2sls-'b0')/(sigmatilda*(abs(phihat)^(-1)*(ssz/N)&(-1/2)))
  - sqrt(N)*'rho0'*sign(phihat)
% }
```