

## Homework 4

### Due Date: November 18 2008

These are all based on problems in Mardia, Kent and Bibby (with minor changes of notation and content): 3.2.6, 3.3.1, 3.4.13, 5.3.3, 5.3.5.

1. Suppose  $(X_1 \ X_2 \ X_3)$  has mean  $(\mu_1 \ \mu_2 \ \mu_3)$  and covariance matrix  $\Sigma = \begin{pmatrix} 1 & \rho & \rho^2 \\ \rho & 1 & 0 \\ \rho^2 & 0 & 1 \end{pmatrix}$ .

Show that the conditional distribution of  $(X_1 \ X_2)$  given  $X_3$  has mean  $(\mu_1 + \rho^2(X_3 - \mu_3) \ \mu_2)$  and covariance matrix  $\begin{pmatrix} 1 - \rho^4 & \rho \\ \rho & 1 \end{pmatrix}$ .

2. If  $X \sim MVN_p(\mu, \Sigma)$  and  $Q\Sigma Q^T$  ( $q \times q$ ) is nonsingular, then given  $QX = q$ , show that the conditional distribution of  $X$  is normal with mean  $\mu + \Sigma Q^T(Q\Sigma Q^T)^{-1}(q - Q\mu)$  and (singular) covariance matrix  $\Sigma - \Sigma Q^T(Q\Sigma Q^T)^{-1}Q\Sigma$ .
3. If  $M \sim W_p(\Sigma, m)$ , show that  $E(M^{-1}) = \frac{\Sigma^{-1}}{m-p-1}$ .
4. (a) Consider the hypothesis  $H_0 : \mu = k\mu_0$  with  $\Sigma$  known (in other words, the hypothesis is that  $\mu$  is some unknown multiple  $k$  of a given vector  $\mu_0$ ). Show that under  $H_0$ , the MLE of  $k$  is  $\hat{k} = \mu_0^T \Sigma^{-1} \bar{X} / \mu_0^T \Sigma^{-1} \mu_0$  where  $\bar{X}$  is the mean of independent  $X_1, \dots, X_n$ , each  $MVN_p(\mu, \Sigma)$ . Also show that the LRT statistic  $\lambda = \frac{L_0}{L_1}$  satisfies

$$-2 \log \lambda = n \bar{X}^T \Sigma^{-1} \{ \Sigma - (\mu_0^T \Sigma^{-1} \mu_0)^{-1} \mu_0 \mu_0^T \} \Sigma^{-1} \bar{X}.$$

Deduce that the *exact* distribution of  $-2 \log \lambda$  is  $\chi_{p-1}^2$  when  $H_0$  is true.

- (b) Now consider the hypothesis  $H_0 : \mu = k\mu_0$  with  $\Sigma$  unknown. In this case the MLE of  $k$  under  $H_0$  is  $\hat{k} = \mu_0^T S_0^{-1} \bar{X} / \mu_0^T S_0^{-1} \mu_0$  (you can assume this without proof). With  $d = \bar{X} - \hat{k}\mu_0$ , show that

$$\begin{aligned} -2 \log \lambda &= n \log(1 + d^T S_0^{-1} d), \\ d^T S_0^{-1} d &= \bar{X}^T S_0^{-1} \{ S_0 - (\mu_0^T S_0^{-1} \mu_0)^{-1} \mu_0 \mu_0^T \} S_0^{-1} \bar{X}. \end{aligned}$$

Hence show that the exact distribution of  $(n-1)d^T S_0^{-1} d$  is  $T_{p-1}^2(n-1)$ .

5. Assume we have a sample of size  $n$  from  $X_i = (X_i^{(1)} \ X_i^{(2)} \ \dots \ X_i^{(p)}) \sim MVN_p(\mu, \Sigma)$ .

- (a) Show that the LRT for the hypothesis that  $X_i^{(1)}$  is uncorrelated with  $(X_i^{(2)} \ \dots \ X_i^{(p)})$  is given by  $\lambda = \frac{L_0}{L_1} = (1 - R^2)^{n/2}$  where  $R^2 = s_{11}^{-1} S_{12} S_{22}^{-1} S_{21}$ ; here  $S = \begin{pmatrix} s_{11} & S_{12} \\ S_{21} & S_{22} \end{pmatrix}$ .

- (b) Consider the null hypothesis that  $\Sigma$  has entries  $\sigma^2$  on the diagonal and  $\rho\sigma^2$  off the diagonal. Let  $S$  be the sample covariance matrix,  $v$  the average of the diagonal entries of  $S$  and  $vr$  the average of the off-diagonal entries. Show that the LRT is

$$\lambda = \left\{ \frac{|S|}{v^p(1-r)^{p-1}(1+(p-1)r)} \right\}^{n/2}.$$

## Some Hints

### Hints on Question 4(b)

The question as stated in Mardia, Kent and Bibby refers to two pieces of theory we have not done in class. Therefore, I am repeating those here.

First, I show the derivation of the formula for  $\hat{k}$  (MKB, page 106): this is actually a bit more involved than just substituting  $S_0$  for  $\Sigma$  in the formula derived in (a). Based on (a) and substituting  $\hat{\Sigma}$ , we have  $\hat{k} = \bar{X}^T \hat{\Sigma}^{-1} \mu_0 / \mu_0^T \hat{\Sigma}^{-1} \mu_0$ . Meanwhile, the equation for  $\hat{\Sigma}$  assuming  $\hat{\mu}$  known is

$$\hat{\Sigma} = S_0 + (\bar{X} - \hat{\mu})(\bar{X} - \hat{\mu})^T \quad (1)$$

(this comes down to minimizing  $\log |\Sigma| + \text{tr}\{S_0 + (\bar{X} - \hat{\mu})(\bar{X} - \hat{\mu})^T\}$  and the solution to that comes from the same argument given in class for the MLE of  $\Sigma$  when  $\mu$  is unconstrained). The problem is to show these two equations are satisfied simultaneously when  $S_0$  is substituted for  $\hat{\Sigma}$  in the formula for  $\hat{k}$ . But if we simultaneously premultiply (1) by  $\hat{\Sigma}^{-1}$  and postmultiply by  $S_0^{-1}$ , then

$$S_0^{-1} = \hat{\Sigma}^{-1} + \hat{\Sigma}^{-1}(\bar{X} - \hat{\mu})(\bar{X} - \hat{\mu})^T S_0^{-1} \quad (2)$$

Now premultiply (2) by  $\mu_0^T$ : we have  $\mu_0^T \hat{\Sigma}^{-1}(\bar{X} - \hat{\mu}) = 0$  from the formula for  $\hat{\mu} = \hat{k} \mu_0$ . Therefore  $\mu_0^T S_0^{-1} = \mu_0^T \hat{\Sigma}^{-1}$ , and it then follows that the two alternative forms of  $\hat{k}$  are the same.

Second, there is a piece of theory for testing a constrained hypothesis about  $\mu$  in the case that  $\Sigma$  is unknown (MKB pp. 132–133). Suppose we want to test the null  $H_0: R\mu = r$  where  $R$  is a  $q \times p$  constraint matrix and  $r$  is given. In this case

$$-2 \log \lambda = n \log(1 + d^T S_0^{-1} d)$$

where  $d = S_0 R^T (R S_0 R^T)^{-1} (R \bar{X} - r)$ . Then

$$(n-1) d^T S_0^{-1} d = (n-1) (R \bar{X} - r)^T (R S_0 R^T)^{-1} (R \bar{X} - r) \quad (3)$$

where  $R \bar{X} \sim MVN_q(r, n^{-1} R \Sigma R^T)$  independently of  $n R S_0 R^T = (n-1) R S R^T \sim W_q(R \Sigma R^T, n-1)$  using Prop. 4 of the notes. Therefore, by definition of Hotelling's  $T^2$ , (3) has distribution  $T_{n-1}^2(q)$ .

### Hints on Question 5

(a) If  $A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}$  then

$$|A| = |A_{11}| \cdot |A_{22} - A_{21} A_{11}^{-1} A_{12}| = |A_{22}| \cdot |A_{11} - A_{12} A_{22}^{-1} A_{21}|.$$

(b) Let  $I$  be the  $p \times p$  identity matrix,  $J$  the  $p \times p$  matrix of ones. If  $E = (1 - \rho)I + \rho J$  is a matrix with 1 on the diagonal and  $\rho$  in all off-diagonal entries, then  $|E| = (1 - \rho)^{p-1} (1 - \rho + p\rho)$  and  $E^{-1} = \frac{1}{1-\rho} \left( I - \frac{\rho}{1-\rho+p\rho} J \right)$ .